

University of Groningen

Role of EPAC in axon determination

Munoz Llancao, Pablo Andrés

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2015

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Munoz Llancao, P. A. (2015). *Role of EPAC in axon determination*. [Thesis fully internal (DIV), University of Groningen]. University of Groningen.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Chapter 5:

A-kinase anchoring proteins: Cyclic AMP compartmentalization in neurodegenerative and obstructive pulmonary diseases

W. J. Poppinga, P. Muñoz-Llancao, C. González-Billault³, M. Schmidt

A-kinase anchoring proteins: Cyclic AMP compartmentalization in neurodegenerative and obstructive pulmonary diseases

W. J. Poppinga^{1,2§}, P. Muñoz-Llancao^{1,3,4}, C. González-Billault³, M. Schmidt^{1,2}

¹University of Groningen, Department of Molecular Pharmacology, Groningen, The Netherlands;

²University of Groningen, University Medical Center Groningen, Groningen Research Institute for Asthma and COPD, GRIAC, Groningen, The Netherlands;

³Laboratory of cell and Neuronal Dynamics (Cenedyn), Department of Biology, Faculty of Sciences, Universidad de Chile;

⁴University of Groningen, University Medical Center Groningen, Department of Neuroscience, Section Medical Physiology, Groningen, The Netherlands.

§Corresponding author

Address:

A. Deusinglaan, 1, 9713 AV

Groningen, The Netherlands

E-mail: w.j.poppinga@rug.nl

Running title: A-kinase anchoring proteins in chronic diseases

Summary

The universal second messenger cyclic AMP (cAMP) is generated upon stimulation of G_s protein-coupled receptors, such as the β_2 -adrenergic receptor (β_2 -AR), and leads to the activation of protein kinase A (PKA), the major cAMP effector protein. PKA oscillates between an on and off state and thereby regulates a plethora of distinct biological responses. The broad activation pattern of PKA and its contribution to several distinct cellular functions lead to the introduction of the concept of compartmentalization of cAMP. A-kinase anchoring proteins (AKAPs) are of central importance due to their unique ability to directly and/or indirectly interact with proteins that either determine the cellular content of cAMP, such as β_2 -AR, adenylyl cyclases and phosphodiesterases, or are regulated by cAMP such as the exchange protein directly activated by cAMP (EPAC). We report on lessons learned from neurons indicating that maintenance of cAMP compartmentalization by AKAP5 is linked to neurotransmission, learning and memory. Disturbance of cAMP compartments seem to be linked to neurodegenerative disease including Alzheimer's disease. We translate this knowledge to compartmentalized cAMP signalling in the lung. Next to AKAP5, we focus here on AKAP12 and Ezrin (AKAP78). These topics will be highlighted in the context of the development of novel pharmacological interventions to tackle AKAP-dependent compartmentalization.

Keywords

cAMP, compartmentalization, AKAP, EPAC

Abbreviations

AC, adenylyl cyclase; AKAP, A-Kinase anchoring protein; AKIP, A-kinase interacting protein; AMPA, 2-amino-3-(3-hydroxy-5-methyl-isoxazol-4-yl)propanoic acid; APP, amyloid precursor protein; β_2 -AR, β_2 -adrenergic receptor; cAMP, cyclic adenosine monophosphate; CREB, cAMP response element-binding protein; CSE, cigarette smoke extract; EPAC, exchange factor directly activated by cAMP; GSKIP, GSK3 β interaction protein; IL-8, interleukin-8; IP₃, inositol 1,4,5-trisphosphate; LTD, long-term depression; LTP, long-term potentiation; MAG, myelin-associated glycoprotein; MAGUK, membrane-associated guanylate kinase; MAP2, microtubule-associated protein 2; MLC, myosin light chain; NFT, neurofibrillary tangle; NMDA, N-Methyl-D-aspartic acid; PDE, phosphodiesterase; PKA, protein kinase A; PKC, protein kinase C; PKI, protein kinase inhibitor; PP2B/CaN, phosphatase 2B/calcineurin (also PPP3); PSD, post-synaptic density; RhoGDI, Rho guanine-nucleotide-dissociation inhibitor; RI, regulatory subunit I of PKA; RII, regulatory subunit II of PKA; SKIP, sphingosine kinase interacting protein.

Introduction

G protein-coupled receptors, such as the G_s -coupled β_2 -adrenergic receptor (β_2 -AR), currently represent one of the largest groups of drug targets (Rask-Andersen *et al.*, 2011). After receptor binding of β_2 -agonists such as isoprenaline and fenoterol, elevation in the cellular content of cyclic adenosine monophosphate (cAMP) is catalysed by membrane-bound adenylyl cyclases (ACs) (Hanoune and Defer, 2001, Beavo and Brunton, 2002, Dessauer, 2009), a process known to be shaped by cAMP-degrading phosphodiesterases (PDEs) (McCahill *et al.*, 2008, Conti and Beavo, 2007, Houslay, 2010, Keravis and Lugnier, 2012, Cheepala *et al.*, 2013). Among the PDE superfamily members, PDE4, PDE7 and PDE8 exhibit substrate specificity towards cAMP (Houslay, 2010, Keravis and Lugnier, 2012).

The best-known effector of cAMP is protein kinase A (PKA). The PKA holoenzyme consists of two catalytic (C) subunits, which exists in three isoforms (C α , C β , C γ), and two regulatory (R) subunits. There are two major isoforms of PKA, designated as PKA(I) and PKA(II), which differ exclusively due to the RI and RII subunits, each again subdivided in an α and β isoform (RI α , RI β , RII α , RII β). Upon binding of two cAMP molecules to each R subunit, the dimer releases the C subunits and thereby initiates target protein phosphorylation. PKA is known to oscillate between an on and off state and thereby regulating a plethora of cellular responses (Taylor *et al.*, 2013). With the discovery of the exchange factor directly activated by cAMP (EPAC) (de Rooij *et al.*, 1998, Kawasaki *et al.*, 1998), the subset of biological functions driven by cAMP started

to become even more diverse (Cheng *et al.*, 2008), and thereby further supported the concept of comPartmentalization of cAMP. Though cyclic nucleotide gated ion channels represent another cAMP targeted group, a detailed description is beyond the scope of our current review and we would like to refer the reader to recent review (Biel, 2009).

Concept of compartmentalization of cAMP

The localisation of the different PKA isoforms and of the EPAC proteins as well as of cAMP generating and degrading enzymes is strictly regulated. Indeed, PKA was already some time ago found activated in either Particulate or soluble cellular fractions (Hayes *et al.*, 1980, Corbin *et al.*, 1977). Clustering of PKA to lipid rafts and caveolae further support the existence of subcellular regions specialised in cAMP signalling that are characterized by a rather dynamic composition of a specific subset of signalling molecules among them G_s-coupled receptors, ACs, PDEs and EPAC (Parton and del Pozo, 2013, Hanzal-Bayer and Hancock, 2007, Gosens *et al.*, 2008, Insel and Ostrom, 2003, Patel *et al.*, 2008).

About 40 years ago, studies primarily performed in heart tissue reported that the two prototypical G_s-coupled receptor agonists isoprenaline and prostaglandin E₁ elevated both the cellular content of cAMP, while only isoprenaline increased cardiac contractility (Hayes *et al.*, 1980, Corbin *et al.*, 1977, Hayes *et al.*, 1979, Buxton and Brunton, 1983). Based on these early studies, the concept of comPartmentalization of cAMP signalling has been introduced and thereby ignited a new surge of cAMP-related research. Since then, several studies provide further insights into the diversity of cellular strategies to compartmentalize intracellular signalling, a concept currently

believed to enable a tightly and fine-tuned control of biological functions. Of particular interest is a recent study from Feinstein and colleagues (Feinstein *et al.*, 2012). Combining mathematical modelling and experimental measurements, the authors demonstrated that the microvascular endothelial barrier strictly rely on subtle local changes in cellular cAMP. Cytosolic produced cAMP disrupted the microvascular endothelial barrier integrity whereas cAMP produced at the plasma membrane increased pulmonary microvascular endothelial barrier integrity (Feinstein *et al.*, 2012). Thus, studies on compartmentalization of cellular cAMP emerged as a theme of central importance to unravel the multiple facets of cAMP signalling and its impact in physiological and pathophysiological settings. Such cAMP gradients may display high spatial resolution as cAMP signalling often occurs within one protein complex orchestrated by a scaffold protein, the most studied family of scaffold proteins coordinating cAMP signalling is the A-Kinase anchoring protein (AKAP) family, outlined in the next paragraph.

A-kinase anchoring proteins

Microtubule-associated protein 2 (MAP2) was the AKAP that tether PKA together with microtubules (Theurkauf and Vallee, 1982). Members of the AKAP family represent important scaffolding proteins and thereby determine the specificity of cellular cAMP signalling. AKAPs control the spatio-temporal activity of the main cAMP effector PKA and some AKAPs have been shown to bind EPAC (Dodge-Kafka *et al.*, 2005, Nijholt *et al.*, 2008, Sehrawat *et al.*, 2011).

Through the association with cAMP-elevating receptors, ACs and/or cAMP-degrading PDEs, AKAPs are able to create and maintain local cAMP pools (Smith *et al.*, 2006) (Smith *et al.*, 2006). To date, over 50 members and splice

variants of the AKAP family have been identified (Skroblin *et al.*, 2010, Welch *et al.*, 2010, Pidoux and Tasken, 2010, Tasken and Aandahl, 2004).

A-kinase anchoring proteins: PKA-RI and PKA-C

Differentiation between AKAPs is based on their ability to bind exclusively PKA-RI, PKA-RII subunits or in the case of dual specific AKAP members both PKA-R subtype. Most of the AKAP superfamily members bind the PKA-RII subunit (Skroblin *et al.*, 2010). In 2010, however, sphingosine kinase interacting protein (SKIP) has been identified as the first mammalian AKAP specific for the binding of PKA-RI (Kovanich *et al.*, 2010, Means *et al.*, 2011, Burgers *et al.*, 2012). In RI α $-/-$ mouse embryonic fibroblasts, SKIP was unable to bind any PKA thereby strongly supporting the notion that SKIP specifically bind PKA-RI (Means *et al.*, 2011). SKIP is also one of the few AKAPs shown to sequester two PKA holoenzymes thereby leading to their sequestration at the inner mitochondrial membrane (Means *et al.*, 2011). Most AKAPs bind with the R subunits and thereby interact also indirectly with the catalytic (C) subunit of PKA. Distinctly different are scaffolding proteins A-kinase interacting protein (AKIP1) (Sastri *et al.*, 2005) and caveolin-1 (Razani *et al.*, 1999) which directly interact with the C subunit. Upon binding to both the C subunit of PKA and the p65 subunit of NF κ B, AKIP1 seems to act as a molecular switch in PKA driven NF κ B signalling (Gao *et al.*, 2010, King *et al.*, 2011). In cardiomyocytes AKIP1 protected against ischemia/reperfusion damage by decreasing reactive oxygen species generation, a process requiring a mitochondrial localization of AKIP1 (Sastri *et al.*, 2013). As both SKIP and AKIP1 seem to exert their primary biological functions in close proximity to mitochondria, it is tempting to speculate

that AKAP scaffolding mechanisms via the PKA-R1 subunit and/or PKA C subunit most likely represent novel molecular mechanisms to unravel yet undefined cellular roles of AKAP-dependent compartmentalization of cAMP.

A-kinase anchoring proteins: Functional diversity and oligomerization

Utilization of distinct combinations of broad-spectrum signalling proteins, such as PKA, protein kinase C (PKC) and protein phosphatase 2B/calcineurin (PP2B/CaN), on the same AKAP, namely AKAP5, modulated the activity of the two distinct neuronal ion channels: 2-amino-3-(3-hydroxy-5-methyl-isoxazol-4-yl)propanoic acid (AMPA)-type glutamate receptor and M-type potassium channels, thereby triggering precise localized cellular responses (Hoshi *et al.*, 2005). With this notion, it is meanwhile generally accepted that AKAPs act as a Swiss army knife that seem to execute differential cellular tasks upon subtle changes in its interacting proteins. Together with the huge number of different members of the AKAP family, the multitude of cellular tasks being performed in different cellular compartments is largely increased.

Even further complexity is added with the finding that AKAPs form homo- (Gao *et al.*, 2011) and hetero-dimers (Gao *et al.*, 2011), a process initially being described for AKAP-Lbc (Baisamy *et al.*, 2005). For example, overexpression of AKAP12 in cells that endogenously express AKAP5, such as HEK293 or A431 cells, potentiates AKAP5-mediated phosphorylation of extracellular signal-regulated kinase 1/2 (ERK1/2) in response to the β_2 -agonist isoprenaline (Gao *et al.*, 2011). Interestingly, however, AKAP12 mediated recycling of the β_2 -AR was unaffected upon AKAP5 overexpression (Gao *et al.*, 2011) (Figure 1). Thus,

oligomerization of AKAP family members may regulate a distinct subset of signalling properties. However, mechanisms involved in AKAP oligomerization, and how such dimer formation is triggered by molecular cues still remain obscure.

For the purpose of this review, we will summarize the most important features of AKAP5, AKAP12, and Ezrin (AKAP78) (Table 1). Neuronal key discoveries will be recapitulated to introduce paradigm shifts that illustrate the general spatio-temporal nature of the compartmentalized cAMP signalling. Our goal is to translate the lessons learned from neurons to the lung as our current knowledge about cAMP compartmentalization in the airways is rather limited. Prior to that, we will focus in the next section on cAMP compartmentalization via AKAPs acting alone with PKA or in concert with EPAC, starting in the following section with the different tools currently available or under development.

Tools to study compartmentalization of cAMP

In the following section, we will highlight novel tools to study the impact of AKAP-bearing multiprotein complexes on a diverse subset of biological functions. As some AKAPs bind next to PKA also EPAC, we will briefly discuss some tools to interfere on the level of PKA or EPAC. For further details about EPAC, we would like to refer the reader to recent reviews on this topic (Oldenburger *et al.*, 2012a, Schmidt *et al.*, 2013, Dekkers *et al.*, 2013). Our main focus are tools to interfere with AKAP-bearing multiprotein complexes.

EPAC and PKA

To distinguish between PKA and EPAC, cell membrane-permeable cyclic nucleotide analogues have been developed, such as N⁶-benzyladenosine-3',5'-cyclic monophosphate (6-Bnz-cAMP) for PKA or 8-(4-chlorophenylthio)-2'-O-methyl-cAMP (8-pCPT-2'-O-Me-cAMP) for EPAC (Schmidt *et al.*, 2013, Holz *et al.*, 2008, Grandoch *et al.*, 2010). In addition, inhibitors of PKA have also been synthesized, such as Rp-8-CPT-cAMPS, and have been shown to abolish the dissociation of PKA-C subunits from the PKA-R subunits (Grandoch *et al.*, 2010). These compounds seem to provide more specificity compared to PKA inhibitors known to act on the ATP binding site, such as H-89 (Bain *et al.*, 2007). Inhibition of PKA can be also achieved with the PKA inhibitor (PKI) (Ibe *et al.*, 2006). Very recently, pharmacological inhibitors of EPAC have been identified, which seem to act primarily on EPAC1 (CE3F4) or EPAC2 (ESI-05) (Tsalkova *et al.*, 2012). Even though researchers world-wide use the novel compounds to gain insight into the contribution of EPAC1 and/or EPAC2 to biological functions (Chen *et al.*, 2013), their mode of action and specificity warrants further studies (Rehmann, 2013). Specific activators for EPAC1 and EPAC2 are still lacking.

A-kinase anchoring proteins: Genetically modified mice

To address the physiological importance of specific AKAPs *in vivo*, mice deficient for a specific AKAP gene e.g. AKAP5^{-/-}, or for a specific AKAP-protein interaction, by introducing a truncation e.g. AKAP5 Δ 36 for AKAP5-PKA interactions, have been developed. Ablation of AKAP members has led to several phenotypes such as decreased fertility, (e.g. AKAP1, AKAP4) cardiac arrhythmias (e.g. AKAP10 (D-AKAP2)), developmental (e.g. mAKAP (AKAP6),

WAVE-1) and neuronal defects (e.g. AKAP5, MAP2) (Hundsrucker and Klusmann, 2008). Based on these findings, it has been suggested that drug targets interfering on the level of AKAPs might bear the ability to disturb signalling driven by cAMP and might therefore represent a novel layer of pharmacological interventions (Dodge-Kafka *et al.*, 2008, Hundsrucker and Klusmann, 2008).

A-kinase anchoring proteins: Dynamics of PKA and AKAP

To assess the dynamics of the primary AKAP interaction partner PKA *in vivo*, several fluorescence resonance energy transfer (FRET) tools have been developed taking advantage of genetically encoded A-kinase activity reporters (Nagai *et al.*, 2000, Zhang *et al.*, 2001, Zhang *et al.*, 2005, Allen and Zhang, 2006, Depry *et al.*, 2011, Komatsu *et al.*, 2011). Addition of cellular localization signals permits the recruitment of these tools to subcellular compartments, including the cytosol, the nucleus, the sarcoplasmic reticulum, mitochondria (using an AKAP based localization), the plasma membrane (Allen and Zhang, 2006, Liu *et al.*, 2011) and even raft or non-raft domains of the cell membrane (Depry *et al.*, 2011). Interestingly, the PKA based biosensors are transferred to the AKAP research upon their combination with AKAP12 (Tao *et al.*, 2010) and AKAP5 (Kocer *et al.*, 2012). Using this novel approach, distinct dynamics of PKA bound to either AKAP12 or AKAP5 at the membrane compared to cytosolic/perinuclear regions were identified (Tao *et al.*, 2010, Kocer *et al.*, 2012). Currently, several novel insights into the subcellular dynamics of AKAP bound PKA are based on cell transduction with PKA defined AKAP reporters and studies in genetically modified mice.

A-kinase anchoring proteins: Pharmacological Tools

The development of a novel pharmacological tool intend to overcome technical limitations and to study the biological impact of AKAP based multiprotein complexes *in vivo*. A conserved amphipathic helix represents a well-defined domain structure present in all AKAP superfamily members, which is required for the interaction with the primary AKAP binding Partner PKA (Skroblin *et al.*, 2010, Malbon *et al.*, 2004, Wang *et al.*, 2006) (Skroblin *et al.*, 2010, Malbon *et al.*, 2004, Wang *et al.*, 2006). The amphipathic helix is inserted into the hydrophobic pockets formed by the dimer of the PKA-R subunits (Kinderman *et al.*, 2006, Gold *et al.*, 2006). It is this amphipathic helix that provided the first basis for the design of dominant interfering peptides able to disrupt the interaction between PKA and AKAP, such as Ht31 (Figure 2A). The stearylated form of Ht31, st-Ht31, exhibits an improved membrane-permeability (Skroblin *et al.*, 2010). It is important to note that the generation of such PKA-AKAP interfering peptides enabled the research community to gain insights into the contribution of AKAP-PKA interactions to a diverse subset of cellular functions in physiology and pathophysiology (Tasken and Aandahl, 2004).

The original peptides, however, provided little, if any, distinction between PKA-RI and PKA-RII subtypes and members of the AKAP family. Through bioinformatics RI (AKB-RI, RIAD) (Burns-Hamuro *et al.*, 2003, Carlson *et al.*, 2006) and RII-specific (AKB-II, (Super)-AKAP-IS) (Burns-Hamuro *et al.*, 2003, Alto *et al.*, 2003) were designed to discriminate between different type of PKA-AKAP interactions, PKA-RI or PKA-RII subunits. In attempts to overcome the central limitation in the current AKAP research field, a recent study from Scott

and colleagues reported on the design of R_{select} peptides, based on the RII subunits of PKA, that seem to exhibit selective affinity for certain members of the AKAP family (Gold *et al.*, 2013). Intriguingly, using phage selection procedure combined with high-resolution structural bioinformatics AKAP2 (AKAP-KL) and AKAP7 (AKAP18) selective R_{select} peptides were validated by biochemical and cell-based experiments (Gold *et al.*, 2013). The AKAP5 (AKAP79, AKAP150) R_{select} peptide, however, did not only interfere with the binding of PKA to AKAP5, but also binding to AKAP7 and AKAP11 (Gold *et al.*, 2013). Functional data for these new tools have yet to come, however, the relevance of this development is evident as for the first time it is possible to distinguish between the individual PKA compartmentalizers without genetic modifications.

In addition, recent studies intend to facilitate a distinction between different AKAPs based on their ability to interact with a discrete interaction partner and/or on mechanisms distinct from the AKAP-PKA interaction outlined above. The dominant interfering peptide GSKIptide, structurally based on the glycogen synthase kinase 3β (GSK3 β) binding site of GSK3 β interaction protein (GSKIP), competes with AKAP members known to bind to GSK3 β , including GSKIP, AKAP11 and MAP2D (in rat) and thereby to disrupt the compartmentalization of GSK3 β (Chou *et al.*, 2006) (Figure 2B). Meanwhile, similar peptides were designed, such as a phospholamban peptide able to prevent the interaction with AKAP7 δ (Lygren *et al.*, 2007) or EBP50 (also known as NHERF1, SLC9A3R1) peptide able to prevent the interaction with Ezrin (AKAP78) (Stokka *et al.*, 2009) (Table 1). Of particular interest are also peptides

that specifically inhibit the interaction between mAKAP and the AC isoform 5 (AC5), leaving the interaction between AKAP5-AC5 unaltered (Kapiloff *et al.*, 2009). Recently, a disruptor for the Hsp20-PDE4 interaction has been described that relief PDE4 from the AKAP-Lbc based complex (Sin *et al.*, 2011).

Most tools being developed thus far, however, are still peptide based and might therefore exert some unknown interactions. For example, it has been reported that st-Ht31P, generated from st-Ht31 by two proline substitution believed to render the molecule in capable of disrupting the AKAP-PKA interaction (Skroblin *et al.*, 2010), seems to inhibit PKA (Klussmann *et al.*, 1999). Current research aims to design small molecule inhibitors for PKA-AKAP interactions (Christian *et al.*, 2011, Schafer *et al.*, 2013). Intriguingly, it has been reported that the small molecule 3,3'-diamino-4,4'-dihydroxydiphenylmethane (FMP-API-1) and its derivatives inhibit AKAP-PKA interactions *in vitro* and in cultured cardiomyocytes (Christian *et al.*, 2011) (Figure 2C). As FMP-API-1, however, also activates PKA (Christian *et al.*, 2011), synthesis of additional small molecules is still warranted. Indeed, new terpyridine scaffolds have been recently synthesized (Schafer *et al.*, 2013), representing the non-peptidic compounds, which might exert less, unwanted biological side effects.

Relation to disease

Disturbance of AKAPs either on the level of their expression profile or biological functions has been associated with a variety of diseases (Tasken and Aandahl, 2004). For example, AKAP12, also known as AKAP250 or Gravin, was first identified as an auto-antigen in myasthenia gravis (Nauert *et al.*, 1997).

Down regulation of AKAP12 is associated with prostate hyperplasia (Akakura *et al.*, 2008) and several types of cancer (Gelman, 2010), including gastric cancer (Choi *et al.*, 2004). It is tempting to speculate that down regulation of AKAP12 might be mediated by promotor hypermethylation, a mechanism described before in the context of oesophageal and colon cancer (Jin *et al.*, 2008, Mori *et al.*, 2006, Paintlia *et al.*, 2009). Such a mechanism is important to promote cancer cell invasiveness by AKAP12 (Su *et al.*, 2010). In line, AKAP12 inhibits cell proliferation (Gelman, 2010, Akakura and Gelman, 2012). Next to AKAP12, other members of the AKAP family such as AKAP4 and AKAP9 are discussed as cancer markers (Chiriva-Internati *et al.*, 2008, Ferrari *et al.*, 2008, Frank *et al.*, 2008, Ferrari *et al.*, 2007, Sharma *et al.*, 2005, Hasegawa *et al.*, 2004).

In the following sections we will first focus on the compartmentalization of cAMP maintained by AKAPs in the context of neuronal learning and memory processes related to neurodegenerative diseases, including Alzheimer's disease, Parkinson's disease, Huntington's disease, multiple sclerosis and Wallerian degeneration. Then, we will highlight our current knowledge about compartmentalized cAMP signalling networks in the context of obstructive pulmonary diseases such as chronic obstructive pulmonary disease and asthma, and whenever appropriate we will emphasize the impact of AKAP-based multiprotein complexes.

Lessons from neurons and neurodegenerative diseases

In the following sections we will discuss the most recent findings on compartmentalized cAMP signalling to maintain proper neuron functions and to

alleviate symptoms of neurodegenerative disease. In particular, we will highlight studies with focus on members of the AKAP family.

Concept of neuronal cAMP compartmentalization: PKA and EPAC

Neurons represent highly polarized structures, displaying short, tapered dendrites and long, thin axons (Andersen and Bi, 2000, Shelly *et al.*, 2010, Hutchins, 2010). In primary rat hippocampal neurons, Poo and colleagues demonstrated that LKB1 phosphorylation by PKA represents an early event in axonal differentiation, whereas Smurf1 phosphorylation by PKA directs selective neuronal degradation of Par6 or RhoA (Shelly *et al.*, 2010, Cheng *et al.*, 2011). In rat dorsal root ganglion neurons, local cAMP levels regulate axonal guidance through attraction and repulsion of axons, a process involving netrin-1 and myelin-associated glycoprotein (MAG) (Murray and Shewan, 2008, Murray *et al.*, 2009). High cAMP levels during the embryonic stage regulate axonal guidance by EPAC, whereas low cAMP levels during the postnatal stage result in growth cone repulsion by PKA (Murray *et al.*, 2009).

Local changes in cAMP determine hippocampus-dependent learning and memory stages such as acquisition, consolidation, retrieval, reconsolidation and extinction (Abel and Nguyen, 2008, Abel and Lattal, 2001). Since the pioneering work in *Aplysia* 30 years ago (Abrams *et al.*, 1984, Castellucci *et al.*, 1980), several studies link cAMP and PKA to locally defined synaptic plasticity, learning and different memory stages (Abel and Nguyen, 2008, Arnsten *et al.*, 2005, Nijholt *et al.*, 2008, Gelinas *et al.*, 2008, Nijholt *et al.*, 2007). In addition, several recent genetic and pharmacological studies report on the role of EPAC in a context-dependent fear-conditioning paradigm (Kelly *et al.*, 2009, Ouyang *et al.*,

2008, Ma *et al.*, 2009, Morozov *et al.*, 2003, Ostroveanu *et al.*, 2010, Schutsky *et al.*, 2011, Srivastava *et al.*, 2012, Yang *et al.*, 2012).

Concept of compartmentalization of cAMP: AKAP5

As outlined above, both PKA and EPAC seem to sense local changes in cAMP to control neuronal development and differentiation, learning and memory. Compartmentalization of cAMP in the brain seems to be maintained primarily by AKAP5 (Moita *et al.*, 2002). AKAP5 is regulated during neuronal development (Robertson *et al.*, 2009), and provides a platform to integrate neuronal cAMP signalling networks (Nijholt *et al.*, 2008). Thus, AKAP5 most likely coordinates the fine tuning of cAMP by regulating the temporal and spatial events controlling cAMP levels. Indeed, a neuronal cAMP-sensing multiprotein complex maintained by AKAP5, PKA, EPAC2 and PKB/Akt, controlled the survival protein kinase B/Akt pathway (Nijholt *et al.*, 2008).

AKAP5: Neurotransmission, learning and memory

Binding of PKA to AKAPs alters synaptic protein phosphorylation and thereby controls synaptic plasticity and memory consolidation (Moita *et al.*, 2002). In hippocampal neurons, AKAP5 acts as a postsynaptic scaffold protein that binds next to PKA protein phosphatase 2B/calcineurin (PP2B/CaN; also PPP3) (Bauman *et al.*, 2004) and PKC (Smith *et al.*, 2006c, Tunquist *et al.*, 2008) (Figure 3). The post-synaptic AKAP5 localization is dependent on its association with the actin cytoskeleton, acidic phospholipids, and cadherins (Gomez *et al.*, 2002, Gorski *et al.*, 2005b). Binding of AKAP5 with membrane-associated guanylate kinase (MAGUK) is required for maturation of dendritic protrusions into large, dendritic spines with an increased density of synaptic AMPA receptors

(Robertson *et al.*, 2009). The functional relation between AKAP5 and AMPA receptors may also be linked to the binding of AKAP5 to the MAGUK family member SAP-97 (Colledge *et al.*, 2000). AKAP5 can also bind the post-synaptic density protein PSD-95 to regulate N-Methyl-D-aspartic acid (NMDA) receptors (Smith *et al.*, 2006b, Bhattacharyya *et al.*, 2009) (Figure 3). Next to the interaction of AKAP5 with several members of the scaffold protein PSD family, binding of AKAP5 to cadherins may also influence synaptic plasticity mechanisms, a process implicated in the regulation of NMDA receptors (Gorski *et al.*, 2005a) (Figure 3).

AKAP5 directly interacts with the neuronal L-type calcium channel subunit Cav1.2 (Oliveria *et al.*, 2007), and thereby forms a complex with AC, PKA and PP2A, capable to modulate Ca^{2+} signalling downstream of the β_2 -AR (Davare *et al.*, 2001). Anchoring of PP2B/CaN to AKAP5 regulates internalization and rapid de-phosphorylation of the AMPA receptor, and most likely reflects a form of molecular and cellular memory associated with long-term depression (LTD) (Figure 3B) (Smith *et al.*, 2006b). Indeed, brain slices derived from adult AKAP5 knock-out mice display normal basal hippocampal spine density and synaptic transmission, but exhibit deficiency in LTD, learning and memory (Robertson *et al.*, 2009). Malenka and colleagues (Jurado *et al.*, 2010) reported that AKAP5 modulates LTD most likely through binding of AKAP5 to PSD-95, causing the release of PP2B/CaN, and subsequently enhances endocytosis of synaptic AMPA receptors. As a consequence, AKAP5 may leave the spine, and thereby contribute to the shrinkage of spines that accompanies LTD (Jurado *et al.*, 2010). Currently, the best genetic models to study AKAP5 function are the $\Delta 36$ mice,

which lack the PKA binding site at the C-terminus of AKAP5 (Lu *et al.*, 2007), and the AKAP5 deficient mice (Weisenhaus *et al.*, 2010). $\Delta 36$ mice display both LTP and LTD defects. In contrast, the AKAP5 deficient mice exhibit only LTD defects. Such differences may suggest that the most critical function of AKAP5 is most likely related to its interaction with PKA, to control the formation and/or maintenance of dendritic spines (Lu *et al.*, 2011). It is clear that regulation of PKA signalling by AKAP5 is necessary to facilitate neurotransmission, learning and defined stages of the memory.

Throughout the mouse brain, AKAP5 is widely distributed in regions linked to learning and memory in rodents, such as the cortex, hippocampus and amygdala (Moita *et al.*, 2002, Glantz *et al.*, 1992, Ostroveanu *et al.*, 2007, Ulfig and Setzer, 1999). Using contextual fear-conditioning in mice, the expression of AKAP5 protein was increased in the hippocampus in a late phase of memory consolidation of associative memory (Nijholt *et al.*, 2007). Disruption of hippocampal AKAP-PKA interactions by st-Ht31 or st-superAKAP-IS facilitates the extinction and impairs consolidation of contextual fear memories, whereas acquisition and retrieval remain unchanged (Nijholt *et al.*, 2008) (Figure 3). Disruption of AKAP-PKA interactions by st-Ht31 in the rat lateral amygdala impaired memory consolidation in auditory fear conditioning (Moita *et al.*, 2002). Using the Morris water maze to study learning and spatial memory, AKAP5 deficient mice exhibit deficits in spatial memory retention most likely caused by delocalization of PKA and subsequent alterations in the local environment of cAMP signalling in the hippocampus (Tunquist *et al.*, 2008). Taken together,

several recent studies *illustrate the importance of AKAP5 to maintain neuronal compartmentalized cAMP signalling to coordinate learning and memory.*

AKAP5: Lessons from Alzheimer's disease

As discussed above, cAMP in neurons is crucial for learning, memory and physiologic events but questions remaining are: How can this system be modulated under pathological neurodegenerative circumstances? Answers most likely provide mechanistic insights that may give some clues for the development of novel pharmacological tools. Ample evidence suggests that perturbation of local cAMP signalling contributes to the development and progression of neurodegenerative diseases. Here we focus on the role of the players discussed previously in the context of Alzheimer's disease.

Alzheimer's disease is a neurodegenerative disease characterized by the progressive decline of cognitive function and memory, and is the fourth largest cause of death for people over 65 years of age (Sonkusare *et al.*, 2005). The disease is characterized by extracellular β -amyloid plaques, intracellular neurofibrillary tangles, cholinergic transmission defects and neuronal loss, preferentially in the entorhinal cortex and hippocampus (Sonkusare *et al.*, 2005). As several inflammatory markers are upregulated in Alzheimer's disease, it is generally assumed that inflammation is linked to the pathogenesis of Alzheimer's disease. Indeed, amyloid plaques seem to trigger inflammatory processes (McGeer and McGeer, 1995, Martinez *et al.*, 1999, Halliday *et al.*, 2000, Rogers, 2008).

Chronic infusion of lipopolysaccharide has been used as experimental model to mimic certain aspects of Alzheimer's disease (Jaeger *et al.*, 2009).

Chronic lipopolysaccharide infusion into the 4th ventricle of young rats induces brain inflammation and subsequently activation of microglial, a process being accompanied by a reduction of adenosine A_{2B} receptor expression and cAMP (Rosi *et al.*, 2003). Mengod and colleagues (Perez-Torres *et al.*, 2003) reported by *in situ* hybridization that at early stages of Alzheimer's disease PDE4, in particular PDE4B, and PDE7 are upregulated, while at later stages of Alzheimer's disease PDE8 is upregulated (Perez-Torres *et al.*, 2003). Both studies imply that progression of Alzheimer's disease is associated with a limitation in cellular cAMP.

Accumulating evidence suggests that A β -induced neurotoxicity alters NMDA receptor signalling through the cAMP response element-binding protein (CREB), a transcription involved in learning and memory processes (Snyder *et al.*, 2005). Moreover, CREB phosphorylation was reduced in the hippocampus of Alzheimer's post-mortem brains (Yamamoto-Sasaki *et al.*, 1999). Intriguingly, Shelanski and colleagues show that treatment of rat hippocampal neurons with A β peptides decreases dissociation of PKA catalytic and regulatory subunits and thereby phosphorylation of downstream targets such as CREB (Vitolo *et al.*, 2002). The PDE4 inhibitor rolipram promotes the dissociation of PKA's catalytic and regulatory subunits and reverses inhibitory effects of A β peptides on CREB phosphorylation (Vitolo *et al.*, 2002, Cheng *et al.*, 2010, Wang *et al.*, 2012). As PKA-dependent signalling studied by CREB phosphorylation in the hippocampus of Alzheimer's post-mortem brains was reduced (Yamamoto-Sasaki *et al.*, 1999), Arima and colleagues proposed that CREB phosphorylation may serve as a

molecular biomarker of ageing-related pathological processes (Sato *et al.*, 2009), in particular of Alzheimer's disease.

Next to PKA, recent studies indicate that EPAC may also be linked to Alzheimer's disease. Lezoualc'h and colleagues show that the EPAC effector Rap1 promotes the activation of Rac, and subsequently leads to the cleavage of the amyloid precursor protein (APP) and production of sAPP α (Maillet *et al.*, 2003). Rap1 can directly interact with STEF, a specific guanine-exchanging factor (GEF) for Rac1, and this association is involved in the secretion of the sAPP α (Zaldua *et al.*, 2007). Moreover, activation of the serotonin receptor of the subtype 4 increases sAPP α through EPAC1/Rap1/Rac (Robert *et al.*, 2005). It has been postulated that sAPP α acts as a memory-enhancer and neuroprotector (Maillet *et al.*, 2003, Robert *et al.*, 2005). Thus, production of sAPP α by EPAC may reduce symptoms of Alzheimer's disease. Indeed, in human brain regions associated with Alzheimer's disease, EPAC1 mRNA is upregulated being accompanied by a down regulation of EPAC2 mRNA (McPhee *et al.*, 2005).

Next to Alzheimer's disease, cAMP and its players are associated with others neurodegenerative disease such as Parkinson's disease, Huntington's disease, multiple sclerosis and Wallerian degeneration (Table 2). Several lines of evidence indicate that alterations of local cAMP dynamics might be caused by inhibition of PKA, upregulation of a specific PDE subset, up/down regulation of EPAC's, or a combination of these events. Persistent limitations in the cellular cAMP level, due to either defect in the cAMP-producing receptors and/or elevations of the cAMP-degrading PDEs, such as PDE4, seem to underpin the development and progression of neurodegenerative diseases (Table 2). Even

though not yet being studied in detail in the context of neurodegenerative diseases, a central role for the AKAP family member AKAP5 might be envisioned due to its ability to interact with the β_2 -AR and/or PDE4 (Lynch *et al.*, 2005), and due to its ability to maintain neuronal cAMP compartmentalization.

Airway smooth muscle and obstructive pulmonary diseases

Chronic obstructive pulmonary disease (COPD) and asthma are both obstructive inflammatory airway diseases characterized by chronic inflammation, airway obstruction and airway remodelling, albeit with different aetiology and specific pathological features (Barnes, 2008, Hogg and Timens, 2009). COPD is predicted to be the third-leading cause of death by disease worldwide in 2020 (Rycroft *et al.*, 2012). Airflow limitation in asthma is reversible with bronchodilators and associated with airway hyperresponsiveness, whereas airway obstruction in COPD is largely irreversible and lung function decline is progressive (Hogg and Timens, 2009, Meurs *et al.*, 2008, Guerra, 2009, Barnes, 2011). Airway smooth muscle cells contribute to disease symptoms in both asthma and COPD due to their multifunctional behavior that supports airway remodeling and airway obstruction, causing the limitation of airflow (Damera and Panettieri, 2011, Halayko *et al.*, 2008, Billington *et al.*, 2013).

Different classes of bronchodilators are used in practice: β_2 -AR agonists (β_2 -agonists), muscarinic receptor antagonists (anticholinergics), individually or in combinations, with or without the addition of anti-inflammatory glucocorticosteroids (Sethi *et al.*, 2012, Kandeel *et al.*, 2013, Peters *et al.*, 2010, Vogelmeier *et al.*, 2011, Meurs *et al.*, 2013). The main targets for the therapeutic treatment of obstructive pulmonary diseases have a direct or indirect link to G

protein-coupled receptor signalling, mainly to the β_2 -AR and the M_3 muscarinic receptor. In obstructive airway diseases, increase in smooth muscle mass and hypercontractility cause severe limitations in the airflow. Airway smooth muscle cell growth is inhibited by several β_2 -agonists such as fenoterol and salbutamol (Ibe *et al.*, 2006). Increased smooth muscle mass is believed to reduce the lumen size of the airways, a process being associated with aberrant β_2 -AR signalling (Deshpande and Penn, 2006). Despite the fact that β_2 -agonists are generally well tolerated (Donohue *et al.*, 2008, Hanania *et al.*, 2010), long term use of β_2 -agonists caused variations in the treatment outcome in asthma and COPD patients, being either less efficacious in COPD patients or even leading to an increased incidence of asthma exacerbations and other markers of morbidity and mortality (Giembycz and Newton, 2006), (Aguilaniu, 2010, Liesker *et al.*, 2002, Kliber *et al.*, 2010).

Another treatment option in obstructive airway diseases is represented by PDE inhibition, for instance the selective PDE4 inhibitors rolipram and roflumilast (Calverley *et al.*, 2009, Rabe, 2011, Global Initiative for Chronic Obstructive Lung Disease, 2010). PDE inhibitors increase the cellular level of cAMP by preventing its degradation. Although both β_2 -agonists and PDE inhibitors show anti-inflammatory properties *in vitro* (Spina, 2008, Kaur *et al.*, 2008, Hallsworth *et al.*, 2001), a notable difference is seen *in vivo*. PDE4 inhibitors show anti-inflammatory properties *in vivo*, but largely lack airway smooth muscle relaxing properties. In contrast, β_2 -agonists show bronchorelaxing properties *in vivo*, but lack anti-inflammatory properties (Calverley *et al.*, 2009, Hurst *et al.*, 2010). Possible explanations for this discrepancy are most likely β_2 -AR desensitization

and/or biased signalling of the β_2 -AR towards ERK signalling (Dickey *et al.*, 2010, Walker *et al.*, 2011), features largely absent by PDE4 inhibitors due to their post-receptor mode of action. Both the process of β_2 -AR desensitization and biased signalling seem to be facilitated by scaffolding proteins such as AKAP5 and AKAP12 (Lefkowitz *et al.*, 2006, Tao and Malbon, 2008) (Figure 1). Subcellular localized cAMP pools seem to cause differential biological effects upon scaffolding protein mediated targeting of either the β_2 -AR or PDEs.

An innovative alternative is, therefore, urgently required to safeguard long term treatment of obstructive lung disorders. Compartmentalized cAMP signalling may provide a novel opportunity for pharmacological interventions. For example, targeting downstream of the β_2 -AR will most likely circumvent receptor desensitization. One might also expect that such strategies will increase treatment specificity, and thereby minimize unwanted side effects, by targeting only the desired cAMP pool. In the following section, the potential impact of compartmentalized cAMP signalling in the lung for further improvement of obstructive airway diseases will be discussed.

A-kinase anchoring proteins: Signalling in the airway smooth muscle

In the airway smooth muscle, the main signalling pathways that determine its functionality are receptors coupling to G_q or G_s proteins. The G_q protein-coupled receptor family is the M_3 muscarinic receptor known to be activated by acetylcholine, and to be inhibited by anticholinergics such as tiotropium (Meurs *et al.*, 2013). After agonist binding, the $G_{\alpha q}$ subunit activates phospholipase C (PLC), thereby leading to the elevation in cellular calcium and activation of calcium/calmodulin-dependent myosin light chain (MLC), a process known to

result in airway smooth muscle contraction (Billington and Penn, 2003, Mizuno and Itoh, 2009). Activation of PKC by diacylglycerol alters also the dephosphorylation of the MLC through several pathways and thereby contributes to the airway smooth muscle tone (Billington and Penn, 2003). Activation of the G_s protein-coupled receptors by drugs targeting the β_2 -AR, causes elevation of cAMP production via G_s and subsequent activation of ACs (Figure 4).

Two members of the AKAP superfamily are known to interact with the β_2 -AR, AKAP5 and AKAP12. Whereas the association of AKAP5 with the β_2 -AR is constitutively (Lynch *et al.*, 2005), agonist binding to the β_2 -AR increases the interaction of the receptor with AKAP12 (Tao *et al.*, 2003). Despite the fact that AKAP5 and AKAP12 share many common features, no redundancy is seen between them with regard to this cellular response (Tao and Malbon, 2008). AKAP5 has been reported to switch the coupling of the β_2 -AR from G_s to G_i , a process most likely facilitated by a PKA-mediated phosphorylation of the receptor (Hill and Baker, 2003, Daaka *et al.*, 1997) (Figure 1). It has been reported that coupling of the β_2 -AR to G_i leads to activation of ERK signalling (Chen and Malbon, 2009). The ERK pathway is known to be link to both proliferative and cytokine production pathways in airway smooth muscle (Roscioni *et al.*, 2011, Roscioni *et al.*, 2011). In the context of obstructive pulmonary diseases, it is worthwhile to emphasize reports indicating that AKAP5 seems to determine the cell surface expression of the β_2 -AR by increasing the affinity of G-protein coupled receptor kinase 2 (GRK2) for $\beta\gamma$ subunits of the G-proteins, causing their translocation to the membrane, leading to the desensitization and internalization of the β_2 -AR (Cong *et al.*, 2001) (Figure 1). In contrast, after desensitization,

AKAP12 is essential for the dephosphorylation, resensitization and recycling of the β_2 -AR back to the cell membrane (Tao and Malbon, 2008, Tao *et al.*, 2003, Chen and Malbon, 2009). In addition, interaction of GRK2 with Ezrin (AKAP78) determines the β_2 -AR internalization (Cant and Pitcher, 2005) (Figure 1).

Based on these findings, it is reasonable to assume that β_2 -AR functions are determined by the balance between AKAP5, AKAP12, and Ezrin (AKAP78) (Figure 1). Indeed, a recent study from Penn and colleagues reported on the expression of AKAP5, AKAP12 and Ezrin (AKAP78) in human airway smooth muscle cells (Horvat *et al.*, 2012). Penn and colleagues did not observe effects of Ht31 or AKAP-*IS* studying whole cell cAMP after stimulation with isoprenaline or the direct AC activator forskolin. However, using a cyclic nucleotide gated ion channel reporter the authors showed that local cAMP concentrations close to the near-membrane compartment were significantly and transiently increased (Horvat *et al.*, 2012). Using a combination of st-Ht31 and a PDE inhibitor cocktail, the authors demonstrated that disruption of PKA-AKAP interactions resulted in sustained AC activity (Horvat *et al.*, 2012). Mathematical models predicted that tethering of PKA to AKAP should cause a three-fold increase of PKA at the β_2 -AR compartment, thereby decreasing input of the β_2 -AR acting as a negative feedback for AC and PDE activity (Horvat *et al.*, 2012). Indeed, direct inhibition of PKA with the protein kinase inhibitor (PKI) completely blunted the rapid decay of the cAMP signal over time (Horvat *et al.*, 2012). With multiple AKAPs possibly involved to create such PKA pool, utilization of tools recently described by Gold *et al.* (Gold *et al.*, 2013) would be necessary to assess the individual contribution of each AKAP.

In the following sections we will discuss the role of cAMP compartmentalization in some of the important features of chronic obstructive pulmonary diseases; contraction, inflammation and remodelling. Herein we will keep the focus on studies performed in airway smooth muscle.

A-kinase anchoring proteins: Airway smooth muscle contraction

Elevation of cAMP leads to the activation of both PKA and EPAC and thereby modulates airway smooth muscle responses (Schmidt *et al.*, 2013, Dekkers *et al.*, 2013). It is well established that PKA on its own deactivates MLC kinase and desensitizes the IP₃ receptor, thereby functionally counteracting the PLC-PKC pathway. In our research group and by others, EPAC has been identified as a novel factor being involved in the regulation of airway smooth muscle relaxation. EPAC, acting most likely via its main effector Rap1, deactivates RhoA and upregulates Rac1 activation, causing the balance to shift from phosphorylated MLC to non-phosphorylated MLC and thus to airway smooth muscle relaxation (Roscioni *et al.*, 2011, Zieba *et al.*, 2011) (Figure 4). Interestingly, Ezrin (AKAP78) is phosphorylated by Rho-regulated Rho-kinase and binds via its ezrin-radixin-moesin domain the Rho inhibitor Rho guanine-nucleotide-dissociation inhibitor (RhoGDI) (Bretscher *et al.*, 2002). Airway smooth muscle cells express both EPAC and Ezrin (AKAP78) (Horvat *et al.*, 2012, Roscioni *et al.*, 2009). Thus deactivation of Rho by EPAC might involve mechanisms driven by Ezrin (AKAP78) and RhoGDI.

In a Madin-Darby canine kidney cell line, activated Ezrin (AKAP78) binds in a calcium dependent manner to Rac and thereby delayed membrane localization of E-cadherin (Pujuguet *et al.*, 2003). Calcium underlies also cellular

compartmentalization and crosstalk with cAMP, a process being facilitated by members of the AKAP family. For example, AKAP5, known to be involved in β_2 -AR desensitization as outlined above (Figure 1), interacts with calcineurin (Coghlan *et al.*, 1995, Oliveria *et al.*, 2003) and calmodulin (Sarkar *et al.*, 1984). Calmodulin competes with PKC in a Ca^{2+} -dependent manner for binding to AKAP5 (Faux and Scott, 1997). More recently, AKAP12, known to be involved in β_2 -AR sensitivity (Figure 1), rapidly redistributes from the plasma membrane to the cytosol upon stimulation with calcium-elevating agents such as ionomycin or thapsigargin (Schott and Grove, 2013). Moreover, it has been reported that AKAP12 displaces PKA-RII from the membrane (Schott and Grove, 2013).

A striking example of cooperativity between cAMP and calcium facilitated by AKAPs is shown for AKAP11 upon assembly of a complex that includes IQGAP1, GSK3 β , and PKA. It has been shown that binding of AKAP11 and IQGAP2 requires high intracellular calcium levels (Logue *et al.*, 2011c, Logue *et al.*, 2011a). At lower intracellular calcium, AKAP11-anchored PKA phosphorylates IQGAP2 and thereby leads to an increase in Rac binding. In the presence of inactive GSK3 β , however, AKAP11 serves as a platform for the assembly of a complex between IQGAP and CLASP2, a plus-end microtubule tracking protein involved in microtubule polymerization. PKA phosphorylation of GSK3 β and elevations in calcium cooperatively drive the formation of an IQGAP1-CLASP2 complex. Both the IQGAP1-Rac and IQGAP1-CLASP2 complexes have been suggested to be involved in microtubule dynamics and cell motility (Logue *et al.*, 2011c, Logue *et al.*, 2011a). AKAP11 was found to be expressed in airway smooth muscle (Horvat *et al.*, 2012). In addition, EPAC not only interacts with

AKAP5, but also with the microtubule network and with the calcium-elevating phospholipase C-epsilon (Schmidt *et al.*, 2013). Future studies should point out if similar mechanisms contribute to airway smooth muscle contraction.

A-kinase anchoring proteins: Airway smooth muscle inflammation

Recently, we reported in human airway smooth muscle cells that direct pharmacological activation of PKA and EPAC synergistically enhances G_q protein-coupled receptor-induced release of the neutrophil chemoattractant interleukin-8 (IL-8) (Roscioni *et al.*, 2009). Silencing of EPAC expression decreased not only IL-8 release in response to EPAC activation but also in response to PKA activation, and *vice versa* PKA inhibition by Rp-8-CPT-cAMPS reduced IL-8 release induced by both PKA and EPAC (Roscioni *et al.*, 2009). Using st-Ht31 to disrupt PKA-AKAP interactions (Figure 2A), preliminary results of our group suggest that PKA and EPAC regulate the IL-8 release in an AKAP dependent manner.

Results from our research groups and others implicate that such close interconnectivity requires the presence of spatial regulation. AKAP5 was shown to be present in the same AKAP-PKA-EPAC complex described before in neuronal cells (Nijholt *et al.*, 2008). In a related study, we showed that induction of IL-8 release by cigarette smoke extract (CSE) was attenuated by the β_2 -agonist fenoterol, seemingly via EPAC and PKA (Oldenburger *et al.*, 2012b). Disturbance of AKAP-based multiprotein complexes might be expected due to the down regulation of EPAC1 and members of the AKAP family by CSE (Oldenburger *et al.*, 2012b, Oldenburger *et al.*, 2014). Indeed, AKAP12 is down regulated in lung cancer (Wikman *et al.*, 2002). With AKAP5 and AKAP12 known

to determine β_2 -AR functions (Figure 1), an important role for PKA and EPAC localization close to G protein-coupled receptors in asthma and COPD could be imagined. This could explain the varying treatment outcomes seen for these bronchodilators in COPD (Aguilaniu, 2010, Kliber *et al.*, 2010, Liesker *et al.*, 2002).

The underlying molecular mechanisms of the attenuation of IL-8 release by cAMP seem to be coordinated via parallel routes. EPAC was shown to inhibit the NF κ B translocation to the nucleus caused by CSE, and PKA counteracts CSE-induced ERK phosphorylation, both known to underlie IL-8 production (Oldenburger *et al.*, 2012b, Saito *et al.*, 2012). Although limited knowledge is currently available on EPAC compartmentalization, both NF κ B and ERK are known to interact with proteins that anchor catalytic and/or regulatory PKA subunits, respectively (Smith *et al.*, 2011, Gao *et al.*, 2008, Gao *et al.*, 2010, King *et al.*, 2011). Thus, it is tempting to speculate that a distinct subset of AKAP members mediate the anti-inflammatory properties of both PKA and EPAC, a research topic open for future investigation.

Our current knowledge implicates that AKAPs are important factors of both inflammation and contraction. The question that remains: what role AKAPs play in airway remodelling?

A-kinase anchoring proteins: Airway smooth muscle remodelling

Another important functional feature of airway smooth muscle cells encompasses the existence of multiple phenotypes, a process reported to involve both PKA and EPAC. Upon chronic exposure to stimuli such as growth factors, ASM cells switch between a contractile and proliferative (synthetic) phenotype

(Halayko *et al.*, 2008). Some researchers have suggested that ASM proliferation is primarily inhibited by EPAC, but not by PKA (Kassel *et al.*, 2008), while others state a more prominent role for PKA (Ibe *et al.*, 2006). Recently, our research group demonstrated that pharmacological activation of either EPAC or PKA prevented platelet-derived growth factor-induced hypocontractility of airway smooth muscle strips and airway smooth muscle proliferation, a process being accompanied by the inhibition of ERK1/2 (Roscioni *et al.*, 2011, Roscioni *et al.*, 2011), suggesting a possible synergism between PKA and EPAC. Our findings were strengthened by other studies in vascular smooth muscle cells (Hewer *et al.*, 2011). Here a concerted action of PKA and EPAC inhibited serum-induced BrdU incorporation, Rb phosphorylation and the expression of cell cycle progression proteins, in a Rap1a-independent fashion (Hewer *et al.*, 2011).

Several signalling pathways have been shown to be involved airway smooth muscle cell proliferation, including ERK1/2 (Lee *et al.*, 2001) and phosphoinositide 3-kinase/Akt (Ibe *et al.*, 2006, Ma *et al.*, 2011). Until now, molecular interactions between the cAMP effectors PKA and EPAC have been studied in great detail in non-pulmonary systems pointing to compartmentalization of both cAMP effectors via muscle specific mAKAP (Dodge-Kafka *et al.*, 2005), via β_2 -AR associated AKAP5 (Nijholt *et al.*, 2008) and via the cytoskeletal scaffolding-AKAP11 complex (Logue *et al.*, 2011). Interestingly, AKAP11 is expressed in ASM using real-time PCR (Horvat *et al.*, 2012). AKAP11 is not only able to bind PKA, but also GSK3, a kinase shown to be involved in expression of contractile proteins in airway smooth muscle (Oenema *et al.*, 2012), their proliferation (Nunes *et al.*, 2008, Gosens *et al.*, 2007)

and pro-fibrotic signalling (Baarsma *et al.*, 2011). Thus, AKAP11 driven cAMP compartmentalization may regulate airway smooth muscle remodelling.

In summary, several lines of evidence point towards the logical conclusion that AKAP family members are most likely of key importance for cAMP compartmentalization and thereby signalling to maintain a fine-tuned control over structural lung cell responses. Future studies will surely add additional insights into our current knowledge of signal compartmentalization and perhaps crosstalk between calcium and cAMP in the lung.

Outlook and Future perspectives

Compartmentalization of cAMP by AKAP family members represents a highly specialized and dynamic process to fine-tune intracellular signalling. Disturbance of cAMP compartmentalization, either due to alterations in AKAP expression or complex composition with a variety of tools outlined herein, seems to profoundly regulate biological functions and thereby to contribute to neurodegenerative and obstructive lung diseases.

Aging of the world-wide population will require further improvement of the management of chronic diseases. Notably, cAMP and its effectors seem to be critical in regulating several processes both in chronic brain and lung diseases. Next to PKA, EPAC seems to act as a novel pharmacological target in both groups of diseases; however, the impact of EPAC compared to PKA might be diverse and sometimes even conflicting. Members of the AKAP superfamily maintain cellular compartmentalization of cAMP primarily via direct interaction with PKA, a process now also linked to EPAC. As AKAP bearing multiprotein complexes regulate receptor desensitization and are able to target simultaneously cAMP and calcium, the AKAP superfamily most likely represent an interesting novel pharmacological concept. In chronic obstructive pulmonary disease, targeting calcium-mediated bronchoconstriction and cAMP-mediated bronchorelaxation by one AKAP related drug might give an additional benefit above the current combination therapy with anticholinergics and β_2 -agonists (Karner and Cates, 2012).

As outlined herein, the design of small molecule inhibitors seems to represent one of the most recent key findings in the field of AKAP research

(Christian *et al.*, 2011, Schafer *et al.*, 2013). The AKAP-PKA interaction has been also used as a template for drug design based on the “Dock-and-Lock Method” (Rossi *et al.*, 2012b, Rossi *et al.*, 2012a). Here, a trivalent drug is created upon conjugation of two identical (pro-)drugs (e.g. interferon-alpha 2b (Rossi *et al.*, 2013)) to the PKA-RII dimer and another drug-(targeting) antibody to an AKAP peptide derived from the amphipathic helix (such as AKAP-IS) (Rossi *et al.*, 2012b, Rossi *et al.*, 2012a, Rossi *et al.*, 2013), a process being stabilized by cysteine residues allowing covalent “locking” of the subunits via disulphide bridges. In theory it should be possible to combine any RII-module with any AKAP-module (Rossi *et al.*, 2012b, Rossi *et al.*, 2012a), the benefit of this method most likely should be envisioned in the creation of a diverse set of potential pharmacological drugs.

Within the AKAP research field, pharmacological tools focus on PKA-AKAP interactions and disruption of other interaction partners from the AKAP complexes. Until now, however, no reports focus on the disruption of AKAP-EPAC complexes. Even though an increasing amount of evidence indicate that EPAC interacts with AKAPs, but also other scaffolds independent of PKA (Schmidt *et al.*, 2013). These Rap-GEF interacting proteins might add another dimension to the concept of subcellular compartmentalization of cAMP, in particular in the context of the physiology and pathophysiology of biological functions.

Acknowledgements

W.J.P. was supported by a grant from the Dutch Lung Foundation (3.2.11.015); P.M.-L. was recipient of an Abel Tasman Talent Program Fellowship from the University of Groningen; C.G.-B. was supported from FONDECYT International, Chile; M.S. was supported by a Rosalind Franklin Fellowship from the University of Groningen and a grant from the Deutsche Forschungsgemeinschaft (IRTG1874/1).

Authors Contributions

Author contributions: W.J.P., P.M.-L., C.G.B., and M.S. conception and design of the review; W.J.P. prepared figures and tables; W.J.P. and M.S. edited and revised the manuscript; M.S. approved final version of manuscript.

Figure legends

Figure 1. Members of the AKAP family and β_2 -AR functioning. Left: AKAP5 has been shown to constitutively associate with the β_2 -AR receptor (Fraser et al., 2000, Lynch *et al.*, 2005, Chen and Malbon, 2009). Upon β_2 -AR activation, AKAP5 bound PKA phosphorylates the receptor, facilitates the switch of G_s to G_i and thereby permits signaling to ERK (Fraser et al., 2000, Lynch *et al.*, 2005). In addition, AKAP5 bound PKA phosphorylates G-protein coupled receptor kinase 2 (GRK2), enhances the affinity of GRK2 for $G_{\beta\gamma}$ subunits and subsequent interaction with the β_2 -AR (Cong *et al.*, 2001). Middle: Receptor bound GRK2 bears the ability to interact with Ezrin (AKAP78), the latter known to be required for the internalization of the β_2 -AR (Cant and Pitcher, 2005). Right: β_2 -AR activation leads also to phosphorylation of AKAP12 via bound PKA and increases the association of AKAP12 with the β_2 -AR receptor, a process known to be essential for the recycling of the β_2 -AR (Tao *et al.*, 2003, Shih *et al.*, 1999).

Figure 2. Strategies to disrupt AKAP complexes. Schematic illustration of the different ways to disrupt AKAP complexes. A, using PKA-AKAP dominant interfering peptides, such as Ht31, to displace PKA as the archetypical AKAP interaction Partner. B, using dominant interfering peptides to disrupt interactions between proteins and AKAPs, such as GSKIptide to remove GSK3 from AKAP complexes. C, Similar strategies are now applied by using small molecules such as FMP-API-1. Further details, see text.

Figure 3. Compartmentalization of cAMP in neurons in relation to

neurodegenerative diseases. A, Illustration of cAMP compartmentalization with emphasize on AKAP5 and selected adaptor proteins in neurons and their alterations under pathological conditions including Alzheimer's disease, Parkinson's disease, Huntington's disease, multiple sclerosis and Wallerian degeneration. B, An example of disrupted cAMP compartmentalization as it was shown that AKAP5 coordinated calcineurin (CaN) was required for AMPA receptor internalization and LTD, removal of the AKAP5 caused an impairment of this LTD. For further details, see text.

Figure 4. Compartmentalization of cAMP in relation to airway smooth

muscle functioning. Schematic illustration of central biological ASM functions, namely contraction, cytokine secretion and proliferation. Endogenous expression of AKAP5, AKAP12 and Ezrin (AKAP78) in ASM seems to maintain defined subcellular signalling compartments. Abbreviations not mentioned in the text; CaM, calmodulin; ROCK, Rho-kinase.

Figures

Figure 1

Fig. 1




Figure 2

Fig. 2




Figure 3

Fig. 3




Figure 4

Fig. 4




Table 1: Subset of AKAP family members known to regulate biological functions in the lung and brain. The most important AKAP interactions are highlighted, except of their primary binding Partner PKA. Text between Parentheses, AKAP synonym using the HUGO gene nomenclature or name of a certain orthologue. For further details and references, see text.

AKAP	Interactions	Processes involved
AKAP5 (HUGO) AKAP79 (Human) AKAP150 (Murine) AKAP75 (Bovine) H21	AKAP12, AKAP5 β_1/β_2 -adrenoceptor ACS, Epac2 PKB/Akt, PKC PP2A/B, Calcineurin, Calmodulin PSD-95, MAGUK, SAP97 PIP ₂ , F-actin E-/N-Cadherin AMPA/NMDA receptor Cav1.2	β_2 -AR switching to ERK β_2 -AR desensitization Cell cycle progression Synaptic plasticity
AKAP12 (HUGO) AKAP250 Gravin (Human) SSeCKS (Murine) Tsga12 Srcs5 AI317366	AKAP5, AKAP12 β_2 -adrenoceptor PKC	β_2 -AR resensitization Cell cycle progression Synaptic plasticity
Ezrin AKAP78 Cytovillin p81 Villin-2	EBP50 (NHERF1) GRK2 RhoGDI Rho Rac Epac	β_2 -AR internalization Actin-binding linker protein

Table 2: cAMP comPartmentalization in neurodegenerative diseases. For further details, see text.

Pathology	Modulator involved	cAMP-dependent effects	References
Alzheimer's disease	PKA	Reduced phosphorylation of CREB	Cheng <i>et al.</i> , 2010
		Inactivation of PKA	Vitolo <i>et al.</i> , 2002
		τ phosphorylation at Ser214 and Ser409	Jicha <i>et al.</i> , 1999
		Down-regulation of A2B receptor/PKA signalling	Rosi <i>et al.</i> , 2003
	Epac	sAPP α production via Epac1/Rap1/Rac	Zaldua <i>et al.</i> , 2007
Parkinson's disease	AKAPs	AKAP79, associated with neurofibrillary pathology	Jicha <i>et al.</i> , 1999
	PDEs	PDE4, PDE4B and PDE7 up-regulation at early stage of Alzheimer's disease	Perez-Torres <i>et al.</i> , 2003
		PDE8 up-regulation at later stage of Alzheimer's disease	
	PKA	Down-regulation of A2A receptor/PKA signalling	Hara <i>et al.</i> , 2010
		α -synuclein stimulates τ phosphorylation by PKA	Qureshi <i>et al.</i> , 2011
Huntington's disease	PDEs	PDE7 and PDE4 inhibition enhances neuroprotection	Morales-Garcia <i>et al.</i> , 2011
			Yang <i>et al.</i> , 2008
	PKA	Decreased levels and CREB activation	Gines <i>et al.</i> , 2003
			Sugars <i>et al.</i> , 2004
Multiple sclerosis	PDEs	Inhibition of PDE4 or PDE10A promotes neuroprotective effects	DeMarch <i>et al.</i> , 2007; Giampa <i>et al.</i> , 2009; 2010
	PKA	β_2 -AR deficient astrocytes produce less cAMP	Chesik <i>et al.</i> , 2008
		Lipoic acid treatment increased PKA activity	Salinthon <i>et al.</i> , 2010
Multiple sclerosis	PDEs	Lovastatin treatment and inhibition of PDE4 promote neuroprotection and neurorepair	Paintlia <i>et al.</i> , 2009

References

- Abel, T, Lattal, KM (2001). Molecular mechanisms of memory acquisition, consolidation and retrieval. *Curr Opin Neurobiol.* 11: 180-7.
- Abel, T, Nguyen, PV (2008). Regulation of hippocampus-dependent memory by cyclic AMP-dependent protein kinase. *Prog Brain Res.* 169: 97-115.
- Abrams, TW, Castellucci, VF, Camardo, JS, Kandel, ER & Lloyd, PE (1984). Two endogenous neuropeptides modulate the gill and siphon withdrawal reflex in *Aplysia* by presynaptic facilitation involving cAMP-dependent closure of a serotonin-sensitive potassium channel. *Proc Natl Acad Sci U S A* 81: 7956-60.
- Aguilaniu, B (2010). Impact of bronchodilator therapy on exercise tolerance in COPD. *International journal of chronic obstructive pulmonary disease* 5: 57-71.
- Akakura, S, Gelman, IH (2012). Pivotal Role of AKAP12 in the Regulation of Cellular Adhesion Dynamics: Control of Cytoskeletal Architecture, Cell Migration, and Mitogenic Signaling. *J Signal Transduct* 2012: 529179, 10.1155/2012/529179.
- Akakura, S, Huang, C, Nelson, PJ, Foster, B & Gelman, IH (2008). Loss of the SSeCKS/Gravin/AKAP12 gene results in prostatic hyperplasia. *Cancer Res* 68: 5096-103, 10.1158/0008-5472.CAN-07-5619.
- Allen, MD, Zhang, J (2006). Subcellular dynamics of protein kinase A activity visualized by FRET-based reporters. *Biochem Biophys Res Commun* 348: 716-21, 10.1016/j.bbrc.2006.07.136.
- Alto, NM, Soderling, SH, Hoshi, N, Langeberg, LK, Fayos, R, Jennings, PA, et al. (2003). Bioinformatic design of A-kinase anchoring protein-in silico: A potent and selective peptide antagonist of type II protein kinase A anchoring. *Proc Natl Acad Sci U S A* 100: 4445-50, DOI 10.1073/pnas.0330734100.
- Andersen, SS, Bi, GQ (2000). Axon formation: a molecular model for the generation of neuronal polarity. *Bioessays* 22: 172-9.
- Arnsten, AF, Ramos, BP, Birnbaum, SG & Taylor, JR (2005). Protein kinase A as a therapeutic target for memory disorders: rationale and challenges. *Trends Mol Med* 11: 121-8.
- Baarsma, HA, Menzen, MH, Halayko, AJ, Meurs, H, Kerstjens, HA & Gosens, R (2011). beta-Catenin signaling is required for TGF-beta1-induced extracellular matrix production by airway smooth muscle cells. *Am J Physiol Lung Cell Mol Physiol* 301: L956-65, 10.1152/ajplung.00123.2011; 10.1152/ajplung.00123.2011.
- Bain, J, Plater, L, Elliott, M, Shpiro, N, Hastie, CJ, McLauchlan, H, et al. (2007). The selectivity of protein kinase inhibitors: a further update. *Biochem J* 408: 297-315, 10.1042/BJ20070797.
- Baisamy, L, Jurisch, N & Diviani, D (2005). Leucine zipper-mediated homo-oligomerization regulates the Rho-GEF activity of AKAP-Lbc. *J Biol Chem* 280: 15405-12, DOI 10.1074/jbc.M414440200.
- Barnes, PJ (2011). Biochemical basis of asthma therapy. *J Biol Chem* 286: 32899-905, 10.1074/jbc.R110.206466; 10.1074/jbc.R110.206466.

- Barnes, PJ (2008). Immunology of asthma and chronic obstructive pulmonary disease. *Nat Rev Immunol* 8: 183-92, 10.1038/nri2254.
- Bauman, AL, Goehring, AS & Scott, JD (2004). Orchestration of synaptic plasticity through AKAP signaling complexes. *Neuropharmacology* 46: 299-310, DOI 10.1016/j.neuropharm.2003.09.016.
- Beavo, JA, Brunton, LL (2002). Cyclic nucleotide research -- still expanding after half a century. *Nat Rev Mol Cell Biol* 3: 710-8, 10.1038/nrm911.
- Bhattacharyya, S, Biou, V, Xu, W, Schluter, O & Malenka, RC (2009). A critical role for PSD-95/AKAP interactions in endocytosis of synaptic AMPA receptors. *Nat Neurosci* 12: 172-81, 10.1038/nn.2249.
- Biel, M (2009). Cyclic nucleotide-regulated cation channels. *J Biol Chem* 284: 9017-21, 10.1074/jbc.R800075200.
- Billington, CK, Ojo, OO, Penn, RB & Ito, S (2013). cAMP regulation of airway smooth muscle function. *Pulm Pharmacol Ther* 26: 112-20, 10.1016/j.pupt.2012.05.007; 10.1016/j.pupt.2012.05.007.
- Billington, CK, Penn, RB (2003). Signaling and regulation of G protein-coupled receptors in airway smooth muscle. *Respir Res* 4: 2.
- Bretscher, A, Edwards, K & Fehon, RG (2002). ERM proteins and merlin: integrators at the cell cortex. *Nat Rev Mol Cell Biol* 3: 586-99, 10.1038/nrm882.
- Burgers, PP, Ma, Y, Margarucci, L, Mackey, M, van der Heyden, MA, Ellisman, M, et al. (2012). A small novel A-kinase anchoring protein (AKAP) that localizes specifically protein kinase A-regulatory subunit I (PKA-RI) to the plasma membrane. *J Biol Chem* 287: 43789-97, 10.1074/jbc.M112.395970; 10.1074/jbc.M112.395970.
- Burns-Hamuro, LL, Ma, Y, Kammerer, S, Reineke, U, Self, C, Cook, C, et al. (2003). Designing isoform-specific peptide disruptors of protein kinase A localization. *Proc Natl Acad Sci U S A* 100: 4072-7, 10.1073/pnas.2628038100 2628038100 [pii].
- Buxton, IL, Brunton, LL (1983). Compartments of cyclic AMP and protein kinase in mammalian cardiomyocytes. *J Biol Chem* 258: 10233-9.
- Calverley, PM, Rabe, KF, Goehring, UM, Kristiansen, S, Fabbri, LM, Martinez, FJ, et al. (2009). Roflumilast in symptomatic chronic obstructive pulmonary disease: two randomised clinical trials. *Lancet* 374: 685-94, 10.1016/S0140-6736(09)61255-1.
- Cant, SH, Pitcher, JA (2005). G protein-coupled receptor kinase 2-mediated phosphorylation of ezrin is required for G protein-coupled receptor-dependent reorganization of the actin cytoskeleton. *Mol Biol Cell* 16: 3088-99, 10.1091/mbc.E04-10-0877.
- Carlson, CR, Lygren, B, Berge, T, Hoshi, N, Wong, W, Tasken, K, et al. (2006). Delineation of type I protein kinase A-selective signaling events using an RI anchoring disruptor. *J Biol Chem* 281: 21535-45, DOI 10.1074/jbc.M603223200.
- Castellucci, VF, Kandel, ER, Schwartz, JH, Wilson, FD, Nairn, AC & Greengard, P (1980). Intracellular injection of the catalytic subunit of cyclic AMP-dependent protein kinase simulates facilitation of transmitter release

- underlying behavioral sensitization in *Aplysia*. *Proc Natl Acad Sci U S A* 77: 7492-6.
- Cheepala, S, Hulot, JS, Morgan, JA, Sassi, Y, Zhang, W, Naren, AP, et al. (2013). Cyclic nucleotide compartmentalization: contributions of phosphodiesterases and ATP-binding cassette transporters. *Annu Rev Pharmacol Toxicol* 53: 231-53, 10.1146/annurev-pharmtox-010611-134609; 10.1146/annurev-pharmtox-010611-134609.
- Chen, H, Wild, C, Zhou, X, Ye, N, Cheng, X & Zhou, J (2013). Recent Advances in the Discovery of Small Molecules Targeting Exchange Proteins Directly Activated by cAMP (EPAC). *J Med Chem*, 10.1021/jm401425e.
- Chen, MH, Malbon, CC (2009). G-protein-coupled receptor-associated A-kinase anchoring proteins AKAP5 and AKAP12: Differential trafficking and distribution. *Cell Signal* 21: 136-42.
- Cheng, PL, Lu, H, Shelly, M, Gao, H & Poo, MM (2011). Phosphorylation of E3 ligase Smurf1 switches its substrate preference in support of axon development. *Neuron* 69: 231-43.
- Cheng, X, Ji, Z, Tsalkova, T & Mei, F (2008). EPAC and PKA: a tale of two intracellular cAMP receptors. *Acta Biochim Biophys Sin (Shanghai)* 40: 651-62.
- Cheng, YF, Wang, C, Lin, HB, Li, YF, Huang, Y, Xu, JP, et al. (2010). Inhibition of phosphodiesterase-4 reverses memory deficits produced by A β 25-35 or A β 1-40 peptide in rats. *Psychopharmacology (Berl)* 212: 181-91.
- Chesik, D, Wilczak, N & De Keyser, J (2008). IGF-1 regulates cAMP levels in astrocytes through a beta2-adrenergic receptor-dependant mechanism. *Int J Med Sci* 5: 240-3.
- Chiriva-Internati, M, Ferrari, R, Yu, Y, Hamrick, C, Gagliano, N, Grizzi, F, et al. (2008). AKAP-4: a novel cancer testis antigen for multiple myeloma. *Br J Haematol* 140: 465-8, DOI 10.1111/j.1365-2141.2007.06940.x.
- Choi, MC, Jong, HS, Kim, TY, Song, SH, Lee, DS, Lee, JW, et al. (2004). AKAP12/Gravin is inactivated by epigenetic mechanism in human gastric carcinoma and shows growth suppressor activity. *Oncogene* 23: 7095-103, 10.1038/sj.onc.1207932.
- Chou, HY, Howng, SL, Cheng, TS, Hsiao, YL, Lieu, AS, Loh, JK, et al. (2006). GSKIP is homologous to the Axin GSK3 β interaction domain and functions as a negative regulator of GSK3 β . *Biochemistry* 45: 11379-89, 10.1021/bi061147r.
- Christian, F, Szaszak, M, Friedl, S, Drewianka, S, Lorenz, D, Goncalves, A, et al. (2011). Small molecule AKAP-protein kinase A (PKA) interaction disruptors that activate PKA interfere with compartmentalized cAMP signaling in cardiac myocytes. *J Biol Chem* 286: 9079-96, 10.1074/jbc.M110.160614; 10.1074/jbc.M110.160614.
- Coghlan, VM, Perrino, BA, Howard, M, Langeberg, LK, Hicks, JB, Gallatin, WM, et al. (1995). Association of Protein-Kinase-a and Protein-Phosphatase-2b with a Common Anchoring Protein. *Science* 267: 108-11.
- Colledge, M, Dean, RA, Scott, GK, Langeberg, LK, Huganir, RL & Scott, JD (2000). Targeting of PKA to glutamate receptors through a MAGUK-AKAP complex. *Neuron* 27: 107-19.

- Cong, M, Perry, SJ, Lin, FT, Fraser, ID, Hu, LA, Chen, W, et al. (2001). Regulation of membrane targeting of the G protein-coupled receptor kinase 2 by protein kinase A and its anchoring protein AKAP79. *J Biol Chem* 276: 15192-9.
- Conti, M, Beavo, J (2007). Biochemistry and physiology of cyclic nucleotide phosphodiesterases: essential components in cyclic nucleotide signaling. *Annu Rev Biochem* 76: 481-511, 10.1146/annurev.biochem.76.060305.150444.
- Corbin, JD, Sugden, PH, Lincoln, TM & Keely, SL (1977). ComPartmentalization of adenosine 3':5'-monophosphate and adenosine 3':5'-monophosphate-dependent protein kinase in heart tissue. *J Biol Chem* 252: 3854-61.
- Courilleau, D, Bissierier, M, Jullian, JC, Lucas, A, Bouyssou, P, Fischmeister, R, et al. (2012). Identification of a tetrahydroquinoline analog as a pharmacological inhibitor of the cAMP-binding protein EPAC. *J Biol Chem* 287: 44192-202, 10.1074/jbc.M112.422956; 10.1074/jbc.M112.422956.
- Daaka, Y, Luttrell, LM & Lefkowitz, RJ (1997). Switching of the coupling of the beta2-adrenergic receptor to different G proteins by protein kinase A. *Nature* 390: 88-91.
- Damera, G, Panettieri, RA, Jr (2011). Does airway smooth muscle express an inflammatory phenotype in asthma? *Br J Pharmacol* 163: 68-80, 10.1111/j.1476-5381.2010.01165.x; 10.1111/j.1476-5381.2010.01165.x.
- Davare, MA, Avdonin, V, Hall, DD, Peden, EM, Burette, A, Weinberg, RJ, et al. (2001). A beta2 adrenergic receptor signaling complex assembled with the Ca²⁺ channel Cav1.2. *Science* 293: 98-101.
- de Rooij, J, Zwartkruis, FJ, Verheijen, MH, Cool, RH, Nijman, SM, Wittinghofer, A, et al. (1998). EPAC is a Rap1 guanine-nucleotide-exchange factor directly activated by cyclic AMP. *Nature* 396: 474-7, 10.1038/24884.
- Dekkers, BG, Racke, K & Schmidt, M (2013). Distinct PKA and EPAC comPartmentalization in airway function and plasticity. *Pharmacol Ther* 137: 248-65, 10.1016/j.pharmthera.2012.10.006; 10.1016/j.pharmthera.2012.10.006.
- DeMarch, Z, Giampa, C, Patassini, S, Martorana, A, Bernardi, G & Fusco, FR (2007). Beneficial effects of rolipram in a quinolinic acid model of striatal excitotoxicity. *Neurobiol Dis* 25: 266-73.
- Depry, C, Allen, MD & Zhang, J (2011). Visualization of PKA activity in plasma membrane microdomains. *Mol Biosyst* 7: 52-8, 10.1039/c0mb00079e; 10.1039/c0mb00079e.
- Deshpande, DA, Penn, RB (2006). Targeting G protein-coupled receptor signaling in asthma. *Cell Signal* 18: 2105-20.
- Dessauer, CW (2009). Adenylyl Cyclase-A-kinase Anchoring Protein Complexes: The Next Dimension in cAMP Signaling. *Mol Pharmacol* 76: 935-41, DOI 10.1124/mol.109.059345.
- Dickey, BF, Walker, JK, Hanania, NA & Bond, RA (2010). beta-Adrenoceptor inverse agonists in asthma. *Curr Opin Pharmacol* 10: 254-9, 10.1016/j.coph.2010.03.002; 10.1016/j.coph.2010.03.002.

- Dodge, KL, Khouangsathiene, S, Kapiloff, MS, Mouton, R, Hill, EV, Houslay, MD, et al. (2001). mAKAP assembles a protein kinase A/PDE4 phosphodiesterase cAMP signaling module. *EMBO J* 20: 1921-30.
- Dodge-Kafka, KL, Bauman, A & Kapiloff, MS (2008). A-kinase anchoring proteins as the basis for cAMP signaling. *Handb Exp Pharmacol*: 3-14, 10.1007/978-3-540-72843-6_1.
- Dodge-Kafka, KL, Souhayer, J, Pare, GC, Carlisle Michel, JJ, Langeberg, LK, Kapiloff, MS, et al. (2005). The protein kinase A anchoring protein mAKAP coordinates two integrated cAMP effector pathways. *Nature* 437: 574-8, nature03966 [pii] 10.1038/nature03966.
- Donohue, JF, Hanania, NA, Sciarappa, KA, Goodwin, E, Grogan, DR, Baumgartner, RA, et al. (2008). Arformoterol and salmeterol in the treatment of chronic obstructive pulmonary disease: a one year evaluation of safety and tolerance. *Ther Adv Respir Dis* 2: 37-48, 10.1177/1753465808089455.
- Faux, MC, Scott, JD (1997). Regulation of the AKAP79-protein kinase C interaction by Ca²⁺/Calmodulin. *J Biol Chem* 272: 17038-44.
- Feinstein, WP, Zhu, B, Leavesley, SJ, Sayner, SL & Rich, TC (2012). Assessment of cellular mechanisms contributing to cAMP compartmentalization in pulmonary microvascular endothelial cells. *Am J Physiol Cell Physiol* 302: C839-52, 10.1152/ajpcell.00361.2011; 10.1152/ajpcell.00361.2011.
- Ferrari, R, Kelly, A, Grizzi, F, Yuefei, Y, Cobos, E & Chiriva-Internati, M (2007). Is AKAP-4 a novel cancer testis antigen for multiple myeloma? *J Invest Med* 55: S266-.
- Ferrari, R, Prabliakar, M, Baglioni, L, Borri, N, Yu, Y, Hyiari, S, et al. (2008). AKAP-4 as biomarker for prostate cancer. *J Invest Med* 56: 394-.
- Frank, B, Wiestler, M, Kropp, S, Hemminki, K, Spurdle, AB, Sutter, C, et al. (2008). Association of a common AKAP9 variant with breast cancer risk: a collaborative analysis. *J Natl Cancer Inst* 100: 437-42, 10.1093/jnci/djn037.
- Fraser, ID, Cong, M, Kim, J, Rollins, EN, Daaka, Y, Lefkowitz, RJ, et al. (2000). Assembly of an A kinase-anchoring protein-beta(2)-adrenergic receptor complex facilitates receptor phosphorylation and signaling. *Curr Biol* 10: 409-12, S0960-9822(00)00419-X [pii].
- Gao, N, Asamitsu, K, Hibi, Y, Ueno, T & Okamoto, T (2008). AKIP1 enhances NF-kappaB-dependent gene expression by promoting the nuclear retention and phosphorylation of p65. *J Biol Chem* 283: 7834-43, 10.1074/jbc.M710285200.
- Gao, N, Hibi, Y, Cueno, M, Asamitsu, K & Okamoto, T (2010). A-kinase-interacting protein 1 (AKIP1) acts as a molecular determinant of PKA in NF-kappaB signaling. *J Biol Chem* 285: 28097-104, 10.1074/jbc.M110.116566.
- Gao, S, Wang, HY & Malbon, CC (2011a). AKAP12 and AKAP5 form higher-order hetero-oligomers. *J Mol Signal* 6: 8, 10.1186/1750-2187-6-8.
- Gao, S, Wang, HY & Malbon, CC (2011b). AKAP5 and AKAP12 Form Homo-oligomers. *J Mol Signal* 6: 3, 10.1186/1750-2187-6-3.
- Gelman, IH (2010). Emerging Roles for SSeCKS/Gravin/AKAP12 in the Control of Cell Proliferation, Cancer Malignancy, and Barrierogenesis. *Genes Cancer* 1: 1147-56, 10.1177/1947601910392984.

- Giampa, C, Laurenti, D, Anzilotti, S, Bernardi, G, Menniti, FS & Fusco, FR (2010). Inhibition of the striatal specific phosphodiesterase PDE10A ameliorates striatal and cortical pathology in R6/2 mouse model of Huntington's disease. *PLoS One* 5: e13417, 10.1371/journal.pone.0013417.
- Giampa, C, Patassini, S, Borreca, A, Laurenti, D, Marullo, F, Bernardi, G, et al. (2009). Phosphodiesterase 10 inhibition reduces striatal excitotoxicity in the quinolinic acid model of Huntington's disease. *Neurobiol Dis* 34: 450-6, 10.1016/j.nbd.2009.02.014.
- Giembycz, MA, Newton, R (2006). Beyond the dogma: novel beta2-adrenoceptor signalling in the airways. *Eur Respir J* 27: 1286-306, 10.1183/09031936.06.00112605.
- Gines, S, Seong, IS, Fossale, E, Ivanova, E, Trettel, F, Gusella, JF, et al. (2003). Specific progressive cAMP reduction implicates energy deficit in presymptomatic Huntington's disease knock-in mice. *Hum Mol Genet* 12: 497-508.
- Glantz, SB, Amat, JA & Rubin, CS (1992). cAMP signaling in neurons: patterns of neuronal expression and intracellular localization for a novel protein, AKAP 150, that anchors the regulatory subunit of cAMP-dependent protein kinase II beta. *Mol Biol Cell* 3: 1215-28.
- Global Initiative for Chronic Obstructive Lung Disease (2010). Global strategy for diagnosis, management, and prevention of COPD (December 2010).
- Gold, MG, Fowler, DM, Means, CK, Pawson, CT, Stephany, JJ, Langeberg, LK, et al. (2013). Engineering AKAP-selective regulatory subunits of PKA through structure-based phage selection. *J Biol Chem*, 10.1074/jbc.M112.447326.
- Gold, MG, Lygren, B, Dokurno, P, Hoshi, N, McConnachie, G, Tasken, K, et al. (2006). Molecular basis of AKAP specificity for PKA regulatory subunits. *Mol Cell* 24: 383-95, DOI 10.1016/j.molcel.2006.09.006.
- Gold, MG, Stengel, F, Nygren, PJ, Weisbrod, CR, Bruce, JE, Robinson, CV, et al. (2011). Architecture and dynamics of an A-kinase anchoring protein 79 (AKAP79) signaling complex. *Proc Natl Acad Sci U S A* 108: 6426-31, 10.1073/pnas.1014400108; 10.1073/pnas.1014400108.
- Gomez, LL, Alam, S, Smith, KE, Horne, E & Dell'Acqua, ML (2002). Regulation of A-kinase anchoring protein 79/150-cAMP-dependent protein kinase postsynaptic targeting by NMDA receptor activation of calcineurin and remodeling of dendritic actin. *J Neurosci* 22: 7027-44.
- Gorski, JA, Gomez, LL, Scott, JD & Dell'Acqua, ML (2005a). Association of an A-kinase-anchoring protein signaling scaffold with cadherin adhesion molecules in neurons and epithelial cells. *Mol Biol Cell* 16: 3574-90.
- Gorski, JA, Gomez, LL, Scott, JD & Dell'Acqua, ML (2005b). Association of an A-kinase-anchoring protein signaling scaffold with cadherin adhesion molecules in neurons and epithelial cells. *Mol Biol Cell* 16: 3574-90, 10.1091/mbc.E05-02-0134.
- Gosens, R, Dueck, G, Rector, E, Nunes, RO, Gerthoffer, WT, Unruh, H, et al. (2007). Cooperative regulation of GSK-3 by muscarinic and PDGF receptors is associated with airway myocyte proliferation. *Am J Physiol Lung Cell Mol Physiol* 293: L1348-58, 10.1152/ajplung.00346.2007.

- Gosens, R, Mutawe, M, Martin, S, Basu, S, Bos, ST, Tran, T, et al. (2008). Caveolae and caveolins in the respiratory system. *Curr Mol Med* 8: 741-53.
- Grandoch, M, Roscioni, SS & Schmidt, M (2010). The role of EPAC proteins, novel cAMP mediators, in the regulation of immune, lung and neuronal function. *Br J Pharmacol* 159: 265-84.
- Guerra, S (2009). Asthma and chronic obstructive pulmonary disease. *Curr Opin Allergy Clin Immunol* 9: 409-16, 10.1097/ACI.0b013e3283300baf; 10.1097/ACI.0b013e3283300baf.
- Halayko, AJ, Tran, T & Gosens, R (2008). Phenotype and functional plasticity of airway smooth muscle: role of caveolae and caveolins. *Proc Am Thorac Soc* 5: 80-8, 10.1513/pats.200705-057VS.
- Halliday, G, Robinson, SR, Shepherd, C & Kril, J (2000). Alzheimer's disease and inflammation: a review of cellular and therapeutic mechanisms. *Clin Exp Pharmacol Physiol* 27: 1-8.
- Hallsworth, MP, Twort, CH, Lee, TH & Hirst, SJ (2001). Beta(2)-Adrenoceptor Agonists Inhibit Release of Eosinophil-Activating Cytokines from Human Airway Smooth Muscle Cells. *Br J Pharmacol* 132: 729-41, 10.1038/sj.bjp.0703866.
- Hanania, NA, Donohue, JF, Nelson, H, Sciarappa, K, Goodwin, E, Baumgartner, RA, et al. (2010). The safety and efficacy of arformoterol and formoterol in COPD. *COPD* 7: 17-31, 10.3109/15412550903499498.
- Hanoune, J, Defer, N (2001). Regulation and role of adenylyl cyclase isoforms. *Annu Rev Pharmacol Toxicol* 41: 145-74, 10.1146/annurev.pharmtox.41.1.145.
- Hanzal-Bayer, MF, Hancock, JF (2007). Lipid rafts and membrane traffic. *FEBS Lett* 581: 2098-104, 10.1016/j.febslet.2007.03.019.
- Hara, M, Fukui, R, Hieda, E, Kuroiwa, M, Bateup, HS, Kano, T, et al. (2010). Role of adrenoceptors in the regulation of dopamine/DARPP-32 signaling in neostriatal neurons. 113: 1046-59.
- Hasegawa, K, Ono, T, Matsushita, H, Shimono, M, Noguchi, Y, Mizutani, Y, et al. (2004). A-kinase anchoring protein 3 messenger RNA expression in ovarian cancer and its implication on prognosis. *International Journal of Cancer* 108: 86-90, Doi 10.1002/Ijc.11565.
- Hayes, JS, Brunton, LL, Brown, JH, Reese, JB & Mayer, SE (1979). Hormonally specific expression of cardiac protein kinase activity. *Proc Natl Acad Sci U S A* 76: 1570-4.
- Hayes, JS, Brunton, LL & Mayer, SE (1980). Selective activation of Particulate cAMP-dependent protein kinase by isoproterenol and prostaglandin E1. *J Biol Chem* 255: 5113-9.
- Hewer, RC, Sala-Newby, GB, Wu, YJ, Newby, AC & Bond, M (2011). PKA and EPAC synergistically inhibit smooth muscle cell proliferation. *J Mol Cell Cardiol* 50: 87-98, 10.1016/j.yjmcc.2010.10.010.
- Hill, SJ, Baker, JG (2003). The ups and downs of Gs- to Gi-protein switching. *Br J Pharmacol* 138: 1188-9.
- Hogg, JC, Timens, W (2009). The pathology of chronic obstructive pulmonary disease. *Annu Rev Pathol* 4: 435-59,

- 10.1146/annurev.pathol.4.110807.092145;
10.1146/annurev.pathol.4.110807.092145.
- Holz, GG, Chepurny, OG & Schwede, F (2008). EPAC-selective cAMP analogs: new tools with which to evaluate the signal transduction properties of cAMP-regulated guanine nucleotide exchange factors. *Cell Signal* 20: 10-20, 10.1016/j.cellsig.2007.07.009.
- Horvat, SJ, Deshpande, DA, Yan, H, Panettieri, RA, Codina, J, DuBose, TD, Jr, et al. (2012). A-kinase anchoring proteins regulate compartmentalized cAMP signaling in airway smooth muscle. *FASEB J* 26: 3670-9, 10.1096/fj.11-201020; 10.1096/fj.11-201020.
- Hoshi, N, Langeberg, LK & Scott, JD (2005). Distinct enzyme combinations in AKAP signalling complexes permit functional diversity. *Nat Cell Biol* 7: 1066-73, Doi 10.1038/Ncb1315.
- Houslay, MD (2010). Underpinning compartmentalised cAMP signalling through targeted cAMP breakdown. *Trends Biochem Sci* 35: 91-100, 10.1016/j.tibs.2009.09.007.
- Hundsrucker, C, Klussmann, E (2008). Direct AKAP-mediated protein-protein interactions as potential drug targets. *Handb Exp Pharmacol*: 483-503, 10.1007/978-3-540-72843-6_20.
- Hurst, JR, Vestbo, J, Anzueto, A, Locantore, N, Mullerova, H, Tal-Singer, R, et al. (2010). Susceptibility to exacerbation in chronic obstructive pulmonary disease. *N Engl J Med* 363: 1128-38, 10.1056/NEJMoa0909883; 10.1056/NEJMoa0909883.
- Hutchins, BI (2010). Competitive outgrowth of neural processes arising from long-distance cAMP signaling. *Sci Signal* 3: jc1.
- Ibe, BO, Portugal, AM & Raj, JU (2006). Levalbuterol inhibits human airway smooth muscle cell proliferation: therapeutic implications in the management of asthma. *Int Arch Allergy Immunol* 139: 225-36, 10.1159/000091168.
- Insel, PA, Ostrom, RS (2003). Forskolin as a tool for examining adenylyl cyclase expression, regulation, and G protein signaling. *Cell Mol Neurobiol* 23: 305-14.
- Jaeger, LB, Dohgu, S, Sultana, R, Lynch, JL, Owen, JB, Erickson, MA, et al. (2009). Lipopolysaccharide alters the blood-brain barrier transport of amyloid beta protein: a mechanism for inflammation in the progression of Alzheimer's disease. *Brain Behav Immun* 23: 507-17.
- Jicha, GA, Weaver, C, Lane, E, Vianna, C, Kress, Y, Rockwood, J, et al. (1999). cAMP-dependent protein kinase phosphorylations on tau in Alzheimer's disease. *J Neurosci* 19: 7486-94.
- Jin, Z, Hamilton, JP, Yang, J, Mori, Y, Olaru, A, Sato, F, et al. (2008). Hypermethylation of the AKAP12 promoter is a biomarker of Barrett's-associated esophageal neoplastic progression. *Cancer Epidemiol Biomarkers Prev* 17: 111-7, 10.1158/1055-9965.EPI-07-0407.
- Jurado, S, Biou, V & Malenka, RC (2010). A calcineurin/AKAP complex is required for NMDA receptor-dependent long-term depression. *Nat Neurosci* 13: 1053-5, 10.1038/nn.2613.
- Kandeel, M, Balaha, M, Inagaki, N & Kitade, Y (2013). Current and future asthma therapies. *Drugs Today (Barc)* 49: 325-39, 10.1358/dot.2013.49.5.1950149; 10.1358/dot.2013.49.5.1950149.

- Kapiloff, MS, Piggott, LA, Sadana, R, Li, J, Heredia, LA, Henson, E, et al. (2009). An adenylyl cyclase-mAKAP β signaling complex regulates cAMP levels in cardiac myocytes. *J Biol Chem* 284: 23540-6, 10.1074/jbc.M109.030072.
- Karner, C, Cates, CJ (2012). Long-acting beta(2)-agonist in addition to tiotropium versus either tiotropium or long-acting beta(2)-agonist alone for chronic obstructive pulmonary disease. *Cochrane Database Syst Rev* 4: CD008989, 10.1002/14651858.CD008989.pub2; 10.1002/14651858.CD008989.pub2.
- Kassel, KM, Wyatt, TA, Panettieri, RA, Jr & Toews, ML (2008). Inhibition of human airway smooth muscle cell proliferation by beta 2-adrenergic receptors and cAMP is PKA independent: evidence for EPAC involvement. *Am J Physiol Lung Cell Mol Physiol* 294: L131-8, 10.1152/ajplung.00381.2007.
- Kaur, M, Holden, NS, Wilson, SM, Sukkar, MB, Chung, KF, Barnes, PJ, et al. (2008). Effect of beta2-adrenoceptor agonists and other cAMP-elevating agents on inflammatory gene expression in human ASM cells: a role for protein kinase A. *Am J Physiol Lung Cell Mol Physiol* 295: L505-14, 10.1152/ajplung.00046.2008.
- Kawasaki, H, Springett, GM, Mochizuki, N, Toki, S, Nakaya, M, Matsuda, M, et al. (1998). A family of cAMP-binding proteins that directly activate Rap1. *Science* 282: 2275-9.
- Kelly, MP, Stein, JM, Vecsey, CG, Favilla, C, Yang, X, Bizily, SF, et al. (2009). Developmental etiology for neuroanatomical and cognitive deficits in mice overexpressing Galphas, a G-protein subunit genetically linked to schizophrenia. *Mol Psychiatry* 14: 398,415, 347.
- Keravis, T, Lugnier, C (2012). Cyclic nucleotide phosphodiesterase (PDE) isozymes as targets of the intracellular signalling network: benefits of PDE inhibitors in various diseases and perspectives for future therapeutic developments. *Br J Pharmacol* 165: 1288-305, 10.1111/j.1476-5381.2011.01729.x; 10.1111/j.1476-5381.2011.01729.x.
- Kinderman, FS, Kim, C, von Daake, S, Ma, Y, Pham, BQ, Spraggon, G, et al. (2006). A dynamic mechanism for AKAP binding to RII isoforms of cAMP-dependent protein kinase. *Mol Cell* 24: 397-408, S1097-2765(06)00663-0 [pii] 10.1016/j.molcel.2006.09.015.
- King, CC, Sastri, M, Chang, P, Pennypacker, J & Taylor, SS (2011). The rate of NF-kappaB nuclear translocation is regulated by PKA and A kinase interacting protein 1. *PLoS One* 6: e18713, 10.1371/journal.pone.0018713.
- Kliber, A, Lynd, LD & Sin, DD (2010). The effects of long-acting bronchodilators on total mortality in patients with stable chronic obstructive pulmonary disease. *Respiratory research* 11: 56.
- Klussmann, E, Maric, K, Wiesner, B, Beyermann, M & Rosenthal, W (1999). Protein kinase A anchoring proteins are required for vasopressin-mediated translocation of aquaporin-2 into cell membranes of renal principal cells. *J Biol Chem* 274: 4934-8.
- Kocer, SS, Wang, HY & Malbon, CC (2012). "Shaping" of cell signaling via AKAP-tethered PDE4D: Probing with AKAR2-AKAP5 biosensor. *J Mol Signal* 7: 4,2187-7-4, 10.1186/1750-2187-7-4; 10.1186/1750-2187-7-4.
- Komatsu, N, Aoki, K, Yamada, M, Yukinaga, H, Fujita, Y, Kamioka, Y, et al. (2011). Development of an optimized backbone of FRET biosensors for

- kinases and GTPases. *Mol Biol Cell* 22: 4647-56, 10.1091/mbc.E11-01-0072; 10.1091/mbc.E11-01-0072.
- Kovanich, D, van der Heyden, MA, Aye, TT, van Veen, TA, Heck, AJ & Scholten, A (2010). Sphingosine kinase interacting protein is an A-kinase anchoring protein specific for type I cAMP-dependent protein kinase. *Chembiochem* 11: 963-71, 10.1002/cbic.201000058.
- Lee, JH, Johnson, PR, Roth, M, Hunt, NH & Black, JL (2001). ERK activation and mitogenesis in human airway smooth muscle cells. *Am J Physiol Lung Cell Mol Physiol* 280: L1019-29.
- Lefkowitz, RJ, Rajagopal, K & Whalen, EJ (2006). New roles for beta-arrestins in cell signaling: not just for seven-transmembrane receptors. *Mol Cell* 24: 643-52, 10.1016/j.molcel.2006.11.007.
- Liesker, JJ, Wijkstra, PJ, Hacken, NHT, Koeter, GH, Postma, DS & Kerstjens, HA (2002). A systematic review of the effects of bronchodilators on exercise capacity in patients with COPD. *Chest* 121: 597-608.
- Liu, S, Zhang, J & Xiang, YK (2011). FRET-based direct detection of dynamic protein kinase A activity on the sarcoplasmic reticulum in cardiomyocytes. *Biochem Biophys Res Commun* 404: 581-6, 10.1016/j.bbrc.2010.11.116; 10.1016/j.bbrc.2010.11.116.
- Logue, JS, Whiting, JL & Scott, JD (2011a). Sequestering Rac with PKA confers cAMP control of cytoskeletal remodeling. *Small GTPases* 2: 173-6, 10.4161/sgtp.2.3.16487.
- Logue, JS, Whiting, JL, Tunquist, B, Langeberg, LK & Scott, JD (2011b). Anchored protein kinase A recruitment of active Rac GTPase. *J Biol Chem* 286: 22113-21, 10.1074/jbc.M111.232660.
- Logue, JS, Whiting, JL, Tunquist, B, Sacks, DB, Langeberg, LK, Wordeman, L, et al. (2011c). AKAP220 protein organizes signaling elements that impact cell migration. *J Biol Chem* 286: 39269-81, 10.1074/jbc.M111.277756; 10.1074/jbc.M111.277756.
- Lu, Y, Lu, YS, Shuai, Y, Feng, C, Tully, T, Xie, Z, et al. (2007). The AKAP Yu is required for olfactory long-term memory formation in *Drosophila*. *Proc Natl Acad Sci U S A* 104: 13792-7.
- Lu, Y, Zha, XM, Kim, EY, Schachtele, S, Dailey, ME, Hall, DD, et al. (2011). A Kinase Anchor Protein 150 (AKAP150)-associated Protein Kinase A Limits Dendritic Spine Density. *J Biol Chem* 286: 26496-506.
- Lygren, B, Carlson, CR, Santamaria, K, Lissandron, V, McSorley, T, Litzenberg, J, et al. (2007). AKAP complex regulates Ca²⁺ re-uptake into heart sarcoplasmic reticulum. *EMBO Rep* 8: 1061-7, DOI 10.1038/sj.embor.7401081.
- Lynch, MJ, Baillie, GS, Mohamed, A, Li, X, Maisonneuve, C, Klussmann, E, et al. (2005). RNA silencing identifies PDE4D5 as the functionally relevant cAMP phosphodiesterase interacting with beta arrestin to control the protein kinase A/AKAP79-mediated switching of the beta(2)-adrenergic receptor to activation of ERK in HEK293B2 cells. *J Biol Chem* 280: 33178-89.
- Ma, L, Brown, M, Kogut, P, Serban, K, Li, X, McConville, J, et al. (2011). Akt activation induces hypertrophy without contractile phenotypic maturation in

- airway smooth muscle. *Am J Physiol Lung Cell Mol Physiol* 300: L701-9, 10.1152/ajplung.00119.2009.
- Ma, N, Abel, T & Hernandez, PJ (2009). Exchange protein activated by cAMP enhances long-term memory formation independent of protein kinase A. *Learn Mem* 16: 367-70, 10.1101/lm.1231009.
- Maillet, M, Robert, SJ, Cacquevel, M, Gastineau, M, Vivien, D, Bertoglio, J, et al. (2003). Crosstalk between Rap1 and Rac regulates secretion of sAPP α . *Nat Cell Biol* 5: 633-9.
- Malbon, CC, Tao, J, Shumay, E & Wang, HY (2004). AKAP (A-kinase anchoring protein) domains: beads of structure-function on the necklace of G-protein signalling. *Biochem Soc Trans* 32: 861-4.
- Martinez, M, Fernandez, E, Frank, A, Guaza, C, de la Fuente, M & Hernanz, A (1999). Increased cerebrospinal fluid cAMP levels in Alzheimer's disease. *Brain Res* 846: 265-7.
- McCahill, AC, Huston, E, Li, X & Houslay, MD (2008). PDE4 associates with different scaffolding proteins: modulating interactions as treatment for certain diseases. *Handb Exp Pharmacol* (186):125-66. doi: 125-66, 10.1007/978-3-540-72843-6_6; 10.1007/978-3-540-72843-6_6.
- McGeer, PL, McGeer, EG (1995). The inflammatory response system of brain: implications for therapy of Alzheimer and other neurodegenerative diseases. *Brain Res Brain Res Rev* 21: 195-218.
- McPhee, I, Gibson, LC, Kewney, J, Darroch, C, Stevens, PA, Spinks, D, et al. (2005). Cyclic nucleotide signalling: a molecular approach to drug discovery for Alzheimer's disease. *Biochem Soc Trans* 33: 1330-2, 10.1042/BST20051330.
- Means, CK, Lygren, B, Langeberg, LK, Jain, A, Dixon, RE, Vega, AL, et al. (2011). An entirely specific type I A-kinase anchoring protein that can sequester two molecules of protein kinase A at mitochondria. *Proc Natl Acad Sci U S A* 108: E1227-35, 10.1073/pnas.1107182108; 10.1073/pnas.1107182108.
- Meurs, H, Gosens, R & Zaagsma, J (2008). Airway hyperresponsiveness in asthma: lessons from in vitro model systems and animal models. *Eur Respir J* 32: 487-502, 10.1183/09031936.00023608; 10.1183/09031936.00023608.
- Meurs, H, Oenema, TA, Kistemaker, LE & Gosens, R (2013). A new perspective on muscarinic receptor antagonism in obstructive airways diseases. *Curr Opin Pharmacol* 13: 316-23, 10.1016/j.coph.2013.04.004; 10.1016/j.coph.2013.04.004.
- Mizuno, N, Itoh, H (2009). Functions and regulatory mechanisms of Gq-signaling pathways. *Neurosignals* 17: 42-54, 10.1159/000186689; 10.1159/000186689.
- Moita, MA, Lamprecht, R, Nader, K & LeDoux, JE (2002). A-kinase anchoring proteins in amygdala are involved in auditory fear memory. *Nat Neurosci* 5: 837-8.
- Morales-Garcia, JA, Redondo, M, Alonso-Gil, S, Gil, C, Perez, C, Martinez, A, et al. (2011). Phosphodiesterase 7 inhibition preserves dopaminergic neurons in cellular and rodent models of Parkinson disease. *PLoS One* 6: e17240.
- Mori, Y, Cai, K, Cheng, Y, Wang, S, Paun, B, Hamilton, JP, et al. (2006). A genome-wide search identifies epigenetic silencing of somatostatin,

- tachykinin-1, and 5 other genes in colon cancer. *Gastroenterology* 131: 797-808, 10.1053/j.gastro.2006.06.006.
- Morozov, A, Muzzio, I, Bourtchouladze, R, Van-Strien, N, Lapidus, K, Yin, D., et al. (2003). Rap1 couples cAMP signaling to a distinct pool of p42/44MAPK regulating excitability, synaptic plasticity, learning, and memory. *Neuron* 39, 309-25.
- Murray, AJ, Shewan, DA (2008). EPAC mediates cyclic AMP-dependent axon growth, guidance and regeneration. *Mol Cell Neurosci* 38: 578-88.
- Murray, AJ, Tucker, SJ & Shewan, DA (2009). cAMP-dependent axon guidance is distinctly regulated by EPAC and protein kinase A. *J Neurosci* 29: 15434-44.
- Nagai, Y, Miyazaki, M, Aoki, R, Zama, T, Inouye, S, Hirose, K, et al. (2000). A fluorescent indicator for visualizing cAMP-induced phosphorylation in vivo. *Nat Biotechnol* 18: 313-6, 10.1038/73767.
- Nauert, JB, Klauck, TM, Langeberg, LK & Scott, JD (1997). Gravin, an autoantigen recognized by serum from myasthenia gravis patients, is a kinase scaffold protein. *Current Biology* 7: 52-62.
- Nijholt, IM, Dolga, AM, Ostroveanu, A, Luiten, PG, Schmidt, M & Eisel, UL (2008). Neuronal AKAP150 coordinates PKA and EPAC-mediated PKB/Akt phosphorylation. *Cell Signal* 20: 1715-24.
- Nijholt, I, Ostroveanu, A, de Bruyn, M, Luiten, P, Eisel, U. & Van der Zee, E. (2007). Both exposure to a novel context and associative learning induce an upregulation of AKAP150 protein in mouse hippocampus. *Neurobiol Learn Mem* 87, 693-6.
- Nunes, RO, Schmidt, M, Dueck, G, Baarsma, H, Halayko, AJ, Kerstjens, HA, et al. (2008). GSK-3/beta-catenin signaling axis in airway smooth muscle: role in mitogenic signaling. *Am J Physiol Lung Cell Mol Physiol* 294: L1110-8, 10.1152/ajplung.00500.2007; 10.1152/ajplung.00500.2007.
- Oenema, TA, Smit, M, Smedinga, L, Racke, K, Halayko, AJ, Meurs, H, et al. (2012). Muscarinic receptor stimulation augments TGF-beta1-induced contractile protein expression by airway smooth muscle cells. *Am J Physiol Lung Cell Mol Physiol* 303: L589-97, 10.1152/ajplung.00400.2011; 10.1152/ajplung.00400.2011.
- Oldenburger, A, Maarsingh, H & Schmidt, M (2012a). Multiple Facets of cAMP Signalling and Physiological Impact: cAMP ComPartmentalization in the Lung. *Pharmaceuticals* 5: 1291-331, 10.3390/ph5121291.
- Oldenburger, A, Roscioni, SS, Jansen, E, Menzen, MH, Halayko, AJ, Timens, W, et al. (2012b). Anti-inflammatory role of the cAMP effectors EPAC and PKA: implications in chronic obstructive pulmonary disease. *PLoS One* 7: e31574, 10.1371/journal.pone.0031574.
- Oldenburger A, Poppinga WJ, Kos F, de Bruin HG, Rijks WF, Heijink IH, et al. (2014). A-kinase anchoring proteins contribute to loss of E-cadherin and bronchial epithelial barrier by cigarette smoke. *Am J Physiol Cell Physiol*. 2014 Mar 15;306(6):C585-97. doi: 10.1152/ajpcell.00183.2013.
- Oliveria, SF, Dell'Acqua, ML & Sather, WA (2007). AKAP79/150 anchoring of calcineurin controls neuronal L-type Ca²⁺ channel activity and nuclear signaling. *Neuron* 55: 261-75.

- Oliveria, SF, Gomez, LL & Dell'Acqua, ML (2003). Imaging kinase-AKAP79-phosphatase scaffold complexes at the plasma membrane in living cells using FRET microscopy. *J Cell Biol* 160: 101-12, DOI 10.1083/jcb.200209127.
- Ostroveanu, A, Van der Zee, EA, Dolga, AM, Luiten, PGM, Eisel, ULM & Nijholt, IM (2007). A-kinase anchoring protein 150 in the mouse brain is concentrated in areas involved in learning and memory. *Brain Res* 1145: 97-107, DOI 10.1016/j.brainres.2007.01.117.
- Ostroveanu, A, van der Zee, E, Eisel, U, Schmidt, M. & Nijholt, I. (2010). Exchange protein activated by cyclic AMP 2 (EPAC2) plays a specific and time-limited role in memory retrieval. *Hippocampus* 20, 1018-26.
- Ouyang, M, Zhang, L, Zhu, JJ, Schwede, F & Thomas, SA (2008). EPAC signaling is required for hippocampus-dependent memory retrieval. *Proc Natl Acad Sci U S A* 105: 11993-7, 0804172105 [pii] 10.1073/pnas.0804172105.
- Paintlia, AS, Paintlia, MK, Singh, I, Skoff, RB & Singh, AK (2009). Combination therapy of lovastatin and rolipram provides neuroprotection and promotes neurorepair in inflammatory demyelination model of multiple sclerosis. *Glia* 57: 182-93, 10.1002/glia.20745.
- Parton, RG, del Pozo, MA (2013). Caveolae as plasma membrane sensors, protectors and organizers. *Nat Rev Mol Cell Biol* 14: 98-112, 10.1038/nrm3512; 10.1038/nrm3512.
- Patel, HH, Murray, F & Insel, PA (2008). G-protein-coupled receptor-signaling components in membrane raft and caveolae microdomains. *Handb Exp Pharmacol* (186):167-84. doi: 167-84, 10.1007/978-3-540-72843-6_7; 10.1007/978-3-540-72843-6_7.
- Perez-Torres, S, Cortes, R, Tolnay, M, Probst, A, Palacios, JM & Mengod, G (2003). Alterations on phosphodiesterase type 7 and 8 isozyme mRNA expression in Alzheimer's disease brains examined by in situ hybridization. *Exp Neurol* 182: 322-34.
- Peters, SP, Kunselman, SJ, Icitovic, N, Moore, WC, Pascual, R, Ameredes, BT, et al. (2010). Tiotropium bromide step-up therapy for adults with uncontrolled asthma. *N Engl J Med* 363: 1715-26, 10.1056/NEJMoa1008770.
- Pidoux, G, Tasken, K (2010). Specificity and spatial dynamics of protein kinase A signaling organized by A-kinase-anchoring proteins. *J Mol Endocrinol* 44: 271-84, 10.1677/JME-10-0010.
- Pujuguet, P, Del Maestro, L, Gautreau, A, Louvard, D & Arpin, M (2003). Ezrin regulates E-cadherin-dependent adherens junction assembly through Rac1 activation. *Mol Biol Cell* 14: 2181-91, 10.1091/mbc.E02-07-0410.
- Qureshi, HY, Paudel, HK (2011). Parkinsonian neurotoxin 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) and alpha-synuclein mutations promote Tau protein phosphorylation at Ser262 and destabilize microtubule cytoskeleton in vitro. 286: 5055-68.
- Rabe, KF (2011). Update on roflumilast, a phosphodiesterase 4 inhibitor for the treatment of chronic obstructive pulmonary disease. *Br J Pharmacol* 163: 53-67, 10.1111/j.1476-5381.2011.01218.x; 10.1111/j.1476-5381.2011.01218.x.
- Rask-Andersen, M, Almen, MS & Schioth, HB (2011). Trends in the exploitation of novel drug targets. *Nat Rev Drug Discov* 10: 579-90, 10.1038/nrd3478; 10.1038/nrd3478.

- Razani, B, Rubin, CS & Lisanti, MP (1999). Regulation of cAMP-mediated signal transduction via interaction of caveolins with the catalytic subunit of protein kinase A. *J Biol Chem* 274: 26353-60.
- Redden, JM, Dodge-Kafka, KL (2011). AKAP phosphatase complexes in the heart. *J Cardiovasc Pharmacol* 58: 354-62, 10.1097/FJC.0b013e31821e5649.
- Rehmann, H (2013). EPAC-inhibitors: facts and artefacts. *Sci Rep* 3: 3032, 10.1038/srep03032; 10.1038/srep03032.
- Robert, S, Mailet, M, Morel, E, Launay, JM, Fischmeister, R, Mercken, L, et al. (2005). Regulation of the amyloid precursor protein ectodomain shedding by the 5-HT₄ receptor and EPAC. *FEBS Lett* 579: 1136-42.
- Robertson, HR, Gibson, ES, Benke, TA & Dell'Acqua, ML (2009). Regulation of postsynaptic structure and function by an A-kinase anchoring protein-membrane-associated guanylate kinase scaffolding complex. *J Neurosci* 29: 7929-43.
- Rogers, J (2008). The inflammatory response in Alzheimer's disease. *J Periodontol* 79: 1535-43, 10.1902/jop.2008.080171; 10.1902/jop.2008.080171.
- Roscioni, SS, Dekkers, BG, Prins, AG, Menzen, MH, Meurs, H, Schmidt, M, et al. (2011). cAMP inhibits modulation of airway smooth muscle phenotype via the exchange protein activated by cAMP (EPAC) and protein kinase A. *Br J Pharmacol* 162: 193-209, 10.1111/j.1476-5381.2010.01011.x; 10.1111/j.1476-5381.2010.01011.x.
- Roscioni, SS, Kistemaker, LE, Menzen, MH, Elzinga, CR, Gosens, R, Halayko, AJ, et al. (2009). PKA and EPAC cooperate to augment bradykinin-induced interleukin-8 release from human airway smooth muscle cells. *Respir Res* 10: 88.
- Roscioni, SS, Maarsingh, H, Elzinga, CR, Schuur, J, Menzen, M, Halayko, AJ, et al. (2011a). EPAC as a novel effector of airway smooth muscle relaxation. *J Cell Mol Med* 15: 1551-63, 10.1111/j.1582-4934.2010.01150.x; 10.1111/j.1582-4934.2010.01150.x.
- Roscioni, SS, Prins, AG, Elzinga, CR, Menzen, MH, Dekkers, BG, Halayko, AJ, et al. (2011b). Functional roles of EPAC and PKA in human airway smooth muscle phenotype plasticity. *Br J Pharmacol*, 10.1111/j.1476-5381.2011.01354.x; 10.1111/j.1476-5381.2011.01354.x.
- Rosi, S, McGann, K, Hauss-Wegrzyniak, B & Wenk, GL (2003). The influence of brain inflammation upon neuronal adenosine A₂B receptors. *J Neurochem* 86: 220-7.
- Rossi, EA, Chang, CH, Cardillo, TM & Goldenberg, DM (2013). Optimization of multivalent bispecific antibodies and immunocytokines with improved in vivo properties. *Bioconjug Chem* 24: 63-71, 10.1021/bc300488f; 10.1021/bc300488f.
- Rossi, EA, Goldenberg, DM & Chang, CH (2012a). Complex and defined biostructures with the dock-and-lock method. *Trends Pharmacol Sci* 33: 474-81, 10.1016/j.tips.2012.06.001; 10.1016/j.tips.2012.06.001.
- Rossi, EA, Goldenberg, DM & Chang, CH (2012b). The dock-and-lock method combines recombinant engineering with site-specific covalent conjugation to generate multifunctional structures. *Bioconjug Chem* 23: 309-23, 10.1021/bc2004999; 10.1021/bc2004999.

- Rycroft, CE, Heyes, A, Lanza, L & Becker, K (2012). Epidemiology of chronic obstructive pulmonary disease: a literature review. *Int J Chron Obstruct Pulmon Dis* 7: 457-94, 10.2147/COPD.S32330; 10.2147/COPD.S32330.
- Saito, T, Sugimoto, N, Ohta, K, Shimizu, T, Ohtani, K, Nakayama, Y, et al. (2012). Phosphodiesterase inhibitors suppress *Lactobacillus casei* cell-wall-induced NF-kappaB and MAPK activations and cell proliferation through protein kinase A--or exchange protein activated by cAMP-dependent signal pathway. *ScientificWorldJournal* 2012: 748572, 10.1100/2012/748572; 10.1100/2012/748572.
- Salinthon, S, Yadav, V, Schillace, RV, Bourdette, DN & Carr, DW (2010). Lipoic acid attenuates inflammation via cAMP and protein kinase A signaling. *PLoS One* 5(9).
- Sarkar, D, Erlichman, J & Rubin, CS (1984). Identification of a calmodulin-binding protein that co-purifies with the regulatory subunit of brain protein kinase II. *J Biol Chem* 259: 9840-6.
- Sastri, M, Barraclough, DM, Carmichael, PT & Taylor, SS (2005). A-kinase-interacting protein localizes protein kinase A in the nucleus. *Proc Natl Acad Sci U S A* 102: 349-54, 10.1073/pnas.0408608102.
- Sastri, M, Haushalter, KJ, Panneerselvam, M, Chang, P, Fridolfsson, H, Finley, JC, et al. (2013). A kinase interacting protein (AKIP1) is a key regulator of cardiac stress. *Proc Natl Acad Sci U S A* 110: E387-96, 10.1073/pnas.1221670110; 10.1073/pnas.1221670110.
- Satoh, J, Tabunoki, H & Arima, K (2009). Molecular network analysis suggests aberrant CREB-mediated gene regulation in the Alzheimer disease hippocampus. *Dis Markers* 27: 239-52, 10.3233/DMA-2009-0670.
- Schafer, G, Milic, J, Eldahshan, A, Gotz, F, Zuhlke, K, Schillinger, C, et al. (2013). Highly Functionalized Terpyridines as Competitive Inhibitors of AKAP-PKA Interactions. *Angew Chem Int Ed Engl*, 10.1002/anie.201304686; 10.1002/anie.201304686.
- Schmidt, M, Dekker, FJ & Maarsingh, H (2013). Exchange protein directly activated by cAMP (EPAC): a multidomain cAMP mediator in the regulation of diverse biological functions. *Pharmacol Rev* 65: 670-709, 10.1124/pr.110.003707; 10.1124/pr.110.003707.
- Schott, M, Grove, B (2013). Receptor-mediated Ca and PKC signaling triggers the loss of cortical PKA compartmentalization through the redistribution of gravin. *Cell Signal*, 10.1016/j.cellsig.2013.07.004; 10.1016/j.cellsig.2013.07.004.
- Schutsky, K, Ouyang, M. & Thomas, S. (2011). Xamoterol impairs hippocampus-dependent emotional memory retrieval via Gi/o-coupled β 2-adrenergic signaling. *Learn Mem* 18, 598-604.
- Scott, JD, Dessauer, CW & Tasken, K (2012). Creating Order from Chaos: Cellular Regulation by Kinase Anchoring. *Annu Rev Pharmacol Toxicol*, 10.1146/annurev-pharmtox-011112-140204.
- Sehrawat, S, Hernandez, T, Cullere, X, Takahashi, M, Ono, Y, Komarova, Y, et al. (2011). AKAP9 regulation of microtubule dynamics promotes EPAC1-induced endothelial barrier properties. *Blood* 117: 708-18, 10.1182/blood-2010-02-268870; 10.1182/blood-2010-02-268870.

- Sethi, S, Mahler, DA, Marcus, P, Owen, CA, Yawn, B & Rennard, S (2012). Inflammation in COPD: implications for management. *Am J Med* 125: 1162-70, 10.1016/j.amjmed.2012.06.024; 10.1016/j.amjmed.2012.06.024.
- Sharma, S, Qian, F, Keitz, B, Driscoll, D, Scanlan, MJ, Skipper, J, et al. (2005). A-kinase anchoring protein 3 messenger RNA expression correlates with poor prognosis in epithelial ovarian cancer. *Gynecol Oncol* 99: 183-8, DOI 10.1016/j.ygyno.2005.06.006.
- Shelly, M, Lim, BK, Cancedda, L, Heilshorn, SC, Gao, H & Poo, MM (2010). Local and long-range reciprocal regulation of cAMP and cGMP in axon/dendrite formation. *Science* 327: 547-52.
- Shih, M, Lin, F, Scott, JD, Wang, HY & Malbon, CC (1999). Dynamic complexes of beta2-adrenergic receptors with protein kinases and phosphatases and the role of gravin. *J Biol Chem* 274: 1588-95.
- Sin, YY, Edwards, HV, Li, X, Day, JP, Christian, F, Dunlop, AJ, et al. (2011). Disruption of the cyclic AMP phosphodiesterase-4 (PDE4)-HSP20 complex attenuates the beta-agonist induced hypertrophic response in cardiac myocytes. *J Mol Cell Cardiol* 50: 872-83, 10.1016/j.yjmcc.2011.02.006.
- Skroblin, P, Grossmann, S, Schafer, G, Rosenthal, W & Klussmann, E (2010). Mechanisms of protein kinase a anchoring. *Int Rev Cell Mol Biol* 283: 235-330, 10.1016/S1937-6448(10)83005-9.
- Smith, FD, Langeberg, LK & Scott, JD (2011). Plugging PKA into ERK scaffolds. *Cell Cycle* 10: 731-2.
- Smith, FD, Langeberg, LK & Scott, JD (2006a). The where's and when's of kinase anchoring. *Trends Biochem Sci* 31: 316-23, DOI 10.1016/j.tibs.2006.04.009.
- Smith, KE, Gibson, ES & Dell'Acqua, ML (2006b). cAMP-dependent protein kinase postsynaptic localization regulated by NMDA receptor activation through translocation of an A-kinase anchoring protein scaffold protein. *J Neurosci* 26: 2391-402.
- Smith, KE, Gibson, ES & Dell'Acqua, ML (2006c). cAMP-dependent protein kinase postsynaptic localization regulated by NMDA receptor activation through translocation of an A-kinase anchoring protein scaffold protein. *J Neurosci* 26: 2391-402, 10.1523/JNEUROSCI.3092-05.2006.
- Snyder, EM, Nong, Y, Almeida, CG, Paul, S, Moran, T, Choi, EY, et al. (2005). Regulation of NMDA receptor trafficking by amyloid-beta. *Nat Neurosci* 8: 1051-8.
- Sonkusare, SK, Kaul, CL & Ramarao, P (2005). Dementia of Alzheimer's disease and other neurodegenerative disorders--memantine, a new hope. *Pharmacol Res* 51: 1-17.
- Spina, D (2008). PDE4 inhibitors: current status. *Br J Pharmacol* 155: 308-15, 10.1038/bjp.2008.307; 10.1038/bjp.2008.307.
- Srivastava, D, Jones, K, Woolfrey, K, Burgdorf, J, Russell, T, Kalmbach, A., et al. (2012). Social, communication, and cortical structural impairments in EPAC2-deficient mice. *J Neurosci* 32, 11864-78.
- Stokka, AJ, Mosenden, R, Ruppelt, A, Lygren, B & Tasken, K (2009). The adaptor protein EBP50 is important for localization of the protein kinase A-

- Ezrin complex in T-cells and the immunomodulating effect of cAMP. *Biochem J* 425: 381-8, 10.1042/BJ20091136.
- Su, B, Bu, Y, Engelberg, D & Gelman, IH (2010). SSeCKS/Gravin/AKAP12 inhibits cancer cell invasiveness and chemotaxis by suppressing a protein kinase C- Raf/MEK/ERK pathway. *J Biol Chem* 285: 4578-86, 10.1074/jbc.M109.073494.
- Sugars, KL, Brown, R, Cook, LJ, Swartz, J & Rubinsztein, DC (2004). Decreased cAMP response element-mediated transcription: an early event in exon 1 and full-length cell models of Huntington's disease that contributes to polyglutamine pathogenesis. *J Biol Chem* 279: 4988-99, 10.1074/jbc.M310226200.
- Tao, J, Malbon, CC (2008). G-protein-coupled receptor-associated A-kinase anchoring proteins AKAP5 and AKAP12: differential signaling to MAPK and GPCR recycling. *J Mol Signal* 3: 19.
- Tao, J, Wang, HY & Malbon, CC (2010). AKAR2-AKAP12 fusion protein "biosenses" dynamic phosphorylation and localization of a GPCR-based scaffold. *J Mol Signal* 5: 3,2187-5-3, 10.1186/1750-2187-5-3; 10.1186/1750-2187-5-3.
- Tao, J, Wang, HY & Malbon, CC (2003). Protein kinase A regulates AKAP250 (gravin) scaffold binding to the beta2-adrenergic receptor. *EMBO J* 22: 6419-29.
- Tasken, K, Aandahl, EM (2004). Localized effects of cAMP mediated by distinct routes of protein kinase A. *Physiol Rev* 84: 137-67, 10.1152/physrev.00021.2003.
- Taylor, SS, Zhang, P, Steichen, JM, Keshwani, MM & Kornev, AP (2013). PKA: Lessons learned after twenty years. *Biochim Biophys Acta* 1834: 1271-8, 10.1016/j.bbapap.2013.03.007; 10.1016/j.bbapap.2013.03.007.
- Theurkauf, WE, Vallee, RB (1982). Molecular characterization of the cAMP-dependent protein kinase bound to microtubule-associated protein 2. *J Biol Chem* 257: 3284-90.
- Troger, J, Moutty, MC, Skroblin, P & Klussmann, E (2012). A-kinase anchoring proteins as potential drug targets. *Br J Pharmacol* 166: 420-33, 10.1111/j.1476-5381.2011.01796.x; 10.1111/j.1476-5381.2011.01796.x.
- Tsalkova, T, Mei, FC & Cheng, X (2012a). A Fluorescence-Based High-Throughput Assay for the Discovery of Exchange Protein Directly Activated by Cyclic AMP (EPAC) Antagonists. *PLoS One* 7: e30441, 10.1371/journal.pone.0030441.
- Tsalkova, T, Mei, FC, Li, S, Chepurny, OG, Leech, CA, Liu, T, et al. (2012b). Isoform-specific antagonists of exchange proteins directly activated by cAMP. *Proc Natl Acad Sci U S A* 109: 18613-8, 10.1073/pnas.1210209109; 10.1073/pnas.1210209109.
- Tunquist, BJ, Hoshi, N, Guire, ES, Zhang, F, Mullendorff, K, Langeberg, LK, et al. (2008). Loss of AKAP150 perturbs distinct neuronal processes in mice. *Proc Natl Acad Sci U S A* 105: 12557-62.
- Ulfing, N, Setzer, M (1999). Expression of a kinase anchoring protein 79 in the human fetal amygdala. *Microsc Res Tech* 46: 48-52.

- Vitolo, OV, Sant'Angelo, A, Costanzo, V, Battaglia, F, Arancio, O & Shelanski, M (2002). Amyloid beta -peptide inhibition of the PKA/CREB pathway and long-term potentiation: reversibility by drugs that enhance cAMP signaling. *Proc Natl Acad Sci U S A* 99: 13217-21.
- Vogelmeier, C, Hederer, B, Glaab, T, Schmidt, H, Rutten-van Molken, MP, Beeh, KM, et al. (2011). Tiotropium versus salmeterol for the prevention of exacerbations of COPD. *N Engl J Med* 364: 1093-103, 10.1056/NEJMoa1008378.
- Walker, JK, Penn, RB, Hanania, NA, Dickey, BF & Bond, RA (2011). New perspectives regarding beta(2) -adrenoceptor ligands in the treatment of asthma. *Br J Pharmacol* 163: 18-28, 10.1111/j.1476-5381.2010.01178.x; 10.1111/j.1476-5381.2010.01178.x.
- Wang, C, Yang, XM, Zhuo, YY, Zhou, H, Lin, HB, Cheng, YF, et al. (2012). The phosphodiesterase-4 inhibitor rolipram reverses Abeta-induced cognitive impairment and neuroinflammatory and apoptotic responses in rats. *Int J Neuropsychopharmacol* 15: 749-66, 10.1017/S1461145711000836; 10.1017/S1461145711000836.
- Wang, HY, Tao, J, Shumay, E & Malbon, CC (2006). G-Protein-coupled receptor-associated A-kinase anchoring proteins: AKAP79 and AKAP250 (gravin). *Eur J Cell Biol* 85: 643-50, S0171-9335(05)00215-3 [pii] 10.1016/j.ejcb.2005.12.003.
- Weisenhaus, M, Allen, ML, Yang, L, Lu, Y, Nichols, CB, Su, T, et al. (2010). Mutations in AKAP5 disrupt dendritic signaling complexes and lead to electrophysiological and behavioral phenotypes in mice. *PLoS One* 5: e10325, 10.1371/journal.pone.0010325.
- Welch, EJ, Jones, BW & Scott, JD (2010). Networking with AKAPs: context-dependent regulation of anchored enzymes. *Mol Interv* 10: 86-97, 10.1124/mi.10.2.6.
- Wikman, H, Kettunen, E, Seppanen, JK, Karjalainen, A, Hollmen, J, Anttila, S, et al. (2002). Identification of differentially expressed genes in pulmonary adenocarcinoma by using cDNA array. *Oncogene* 21: 5804-13.
- Willoughby, D, Masada, N, Wachten, S, Pagano, M, Halls, ML, Everett, KL, et al. (2010). A-kinase anchoring protein 79/150 interacts with adenylyl cyclase type 8 and regulates Ca²⁺-dependent cAMP synthesis in pancreatic and neuronal systems. *J Biol Chem*, 10.1074/jbc.M110.120725.
- Yamamoto-Sasaki, M, Ozawa, H, Saito, T, Rosler, M & Riederer, P (1999). Impaired phosphorylation of cyclic AMP response element binding protein in the hippocampus of dementia of the Alzheimer type. 824: 300-3.
- Yan, H, Deshpande, DA, Misior, AM, Miles, MC, Saxena, H, Riemer, EC, et al. (2011). Anti-mitogenic effects of beta-agonists and PGE2 on airway smooth muscle are PKA dependent. *FASEB J* 25: 389-97, 10.1096/fj.10-164798.
- Yang, L, Calingasan, NY, Lorenzo, BJ & Beal, MF (2008). Attenuation of MPTP neurotoxicity by rolipram, a specific inhibitor of phosphodiesterase IV. *Exp Neurol* 211: 311-4, 10.1016/j.expneurol.2007.02.010.
- Yang, Y, Shu, X, Liu, D, Shang, Y, Wu, Y, Pei, L., et al. (2012). EPAC null mutation impairs learning and social interactions via aberrant regulation of miR-124 and Zif268 translation. *Neuron* 73, 774-88.

- Zaldua, N, Gastineau, M, Hoshino, M, Lezoualc'h, F & Zugaza, JL (2007). EPAC signaling pathway involves STEF, a guanine nucleotide exchange factor for Rac, to regulate APP processing. *FEBS Lett* 581: 5814-8.
- Zhang, J, Hupfeld, CJ, Taylor, SS, Olefsky, JM & Tsien, RY (2005). Insulin disrupts beta-adrenergic signalling to protein kinase A in adipocytes. *Nature* 437: 569-73, 10.1038/nature04140.
- Zhang, J, Ma, Y, Taylor, SS & Tsien, RY (2001). Genetically encoded reporters of protein kinase A activity reveal impact of substrate tethering. *Proc Natl Acad Sci U S A* 98: 14997-5002, 10.1073/pnas.211566798.
- Zieba, BJ, Artamonov, MV, Jin, L, Momotani, K, Ho, R, Franke, AS, et al. (2011). The cAMP-responsive Rap1 guanine nucleotide exchange factor, EPAC, induces smooth muscle relaxation by down-regulation of RhoA activity. *J Biol Chem* 286: 16681-92, 10.1074/jbc.M110.205062.